

***Vibrio cholerae*, Strain PR542**

Catalog No. NR-155

(Derived from ATCC® 14103™)

For research use only. Not for human use.

Contributor:

ATCC®

Product Description:

Bacteria Classification: *Vibrionaceae*, *Vibrio*

Species: *Vibrio cholerae*

Strain: PR542

Biotype: Classical

Serogroup: O:1

Bacteriophage Type: Group IV (S. Mukerjee)^{1,2}

Original Source: Human clinical specimen from the cholera epidemic in Calcutta, India (circa 1960)

Comment: *Vibrio cholerae* (*V. cholerae*), strain PR542 was deposited at ATCC® in 1960 by Dr. S. Mukerjee, Division of Microbiology, Indian Institute of Biochemistry and Experimental Medicine, Calcutta, India.

V. cholerae is a natural inhabitant of warm aquatic environments and the causative agent of the diarrheal disease cholera. More than 200 O-antigen serogroups have been identified but only O1 and more recently O139 are known to cause epidemic and pandemic cholera.³ Occasionally, there are cholera outbreaks that result from non-O1 and non-O139 serotypes. *V. cholerae* colonizes the mucosal surface of the small intestines of humans, the only known animal host.⁴ Cholera has a high lethality if left untreated, and millions have died in the seven pandemics that have occurred since 1817.

Cholera toxin, the toxin-coregulated pilus (TCP) and the central regulatory protein, ToxR, are recognized as significant factors in the pathogenicity of toxigenic strains of *V. cholerae* serogroups O1 and O139.⁵

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in 0.5X Tryptic Soy Broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-155 was packaged aseptically in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy Broth or equivalent

Tryptic Soy Agar or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Vibrio cholerae*, Strain PR542, NR-155."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmbl5/bmbl5toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Mukerjee, S., D. K. Guha, and U. K. Guha Roy. "Studies on Typing of Cholera by Bacteriophage. I. Phage-Typing of *Vibrio cholerae* from Calcutta Epidemics." Ann. Biochem. Exp. Med. 17 (1957): 161-176.
2. Basu, S. and S. Mukerjee. "Bacteriophage Typing of *Vibrio* El Tor." Experientia 24 (1968): 299-300. PubMed: 5661438.
3. Pang, B., et al. "Genetic Diversity of Toxigenic and Nontoxigenic *Vibrio cholerae* Serogroups O1 and O139 Revealed by Array-Based Comparative Genomic Hybridization." J. Bacteriol. 189 (2007): 4837-4849. PubMed: 17468246.
4. O'Shea, Y. A., et al. "Evolutionary Genetic Analysis of the Emergence of Epidemic *Vibrio cholerae* Isolates on the Basis of Comparative Nucleotide Sequence Analysis and Multilocus Virulence Gene Profiles." J. Clin. Microbiol. 42 (2004): 4657-4671. PubMed: 15472325.
5. Singh, D. V., et al. "Molecular Analysis of *Vibrio cholerae* O1, O139, non-O1, and non-O139 Strains: Clonal Relationships between Clinical and Environmental Isolates." Appl. Environ. Microbiol. 67 (2001): 910-921. PubMed: 11157262.
6. Mukerjee, S. "Bacteriophage Typing of Cholera." Bull. World Health Org. 28 (1963): 337-345.

ATCC® is a trademark of the American Type Culture Collection.

