

**Toxoplasma gondii, CTG.11 ARA-SNF**

**Catalog No. NR-10273**

**Product Description:** *Toxoplasma gondii* (*T. gondii*), CTG.11 ARA-SNF was the product of a genetic cross between singly resistant parental clones of the C (also CEP and CTG) strain, that were obtained by chemical mutagenesis.

**Lot<sup>1</sup>: 58499196**

**Manufacturing Date: 10APR2009**

TEST	SPECIFICATIONS	RESULTS
<b>Genotyping<sup>2,3</sup></b> AK16 locus ( <i>Hinf</i> I digestion) L358 locus ( <i>Hae</i> III digestion)	Consistent with parental Type III strain Consistent with parental Type III strain	Consistent with parental Type III strain Consistent with parental Type III strain
<b>Drug susceptibility<sup>4</sup></b> Sinefungin (SNF) Adenine arabinoside (Ara-A)	Resistant Resistant	Resistant Resistant
<b>Viable Cell Count by Hemacytometry (pre-freeze)</b>	> 10 <sup>6</sup> cells/mL	2.4 x 10 <sup>7</sup> cells/mL
<b>Viability (post-freeze)<sup>5</sup></b>	Growth	Growth
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Brain Heart Infusion, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> DNA Detection by PCR	None detected	None detected

<sup>1</sup>NR-10273 was produced by cultivation of the deposited material in human foreskin fibroblast cells (ATCC<sup>®</sup> CRL-1634<sup>™</sup>) with cell cultivation medium for parasites ([ATCC medium 2222](#); adjusted to contain 10% heat-inactivated fetal bovine serum). The culture was propagated in 95% air, 5% CO<sub>2</sub> for 10 days at 37°C, until lysis of the host cell monolayer was reached.

<sup>2</sup>PCR amplification was performed separately for the two loci AK16 and L358. Where appropriate, samples were subjected to restriction enzyme digestion typing by agarose gel electrophoresis.

<sup>3</sup>Primer sequences, annealing temperatures, and conditions for restriction enzyme digestion may be obtained at the *Toxoplasma* Genome Map website ([Toxoplasma Genome Map](#)).

<sup>4</sup>Sinefungin was used at a concentration of 2.7 x 10<sup>-7</sup> M and ara-A was used at a concentration of 1.3 x 10<sup>-4</sup> M, as described (Sibley, L. D., et al. "Generation of a Restriction Fragment Length Polymorphism Linkage Map for *Toxoplasma gondii*." *Genetics* 132 (1992): 1003-1015. PubMed: 1360931.)

<sup>5</sup>Viable cells and signs of infection were seen after 10 days under cultivation conditions at 37°C.

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Date: 31 AUG 2009**

**Signature: Signature on File**

**Title: Technical Manager, BEI Authentication or designee**

ATCC<sup>®</sup>, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC<sup>®</sup>'s knowledge.

ATCC<sup>®</sup> is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

