b|**e**|**i** resources

Brugia malayi Microfilariae from the Peritoneal Cavity of a Mongolian Gerbil (Frozen)

Catalog No. NR-48889

This reagent is the tangible property of the U.S. Government.

For research use only. Not for human use.

Contributor:

Andrew R. Moorhead, D.V.M., M.S., Ph.D., Director and Principal Investigator, Filariasis Research Reagent Resource Center, Department of Infectious Diseases University of Georgia College of Veterinary Medicine, Athens, Georgia, USA

Manufacturer:

Filariasis Research Reagent Resource Center supported by Contract HHSN272201000030I, NIH-NIAID Animal Models of Infectious Disease Program

Product Description:

<u>Classification</u>: Onchocercidae, Brugia <u>Species</u>: Brugia malayi <u>Strain</u>: FR3

<u>Original Source:</u> *Brugia malayi* (*B. malayi*), strain FR3 was originally obtained from researchers in Malaysia by Dr. John Schacher.^{1,2}

B. malayi is a roundworm nematode and one of the three causative agents of lymphatic filariasis in humans.³ Lymphatic filariasis, also known as elephantiasis, is a condition characterized by swelling of the lower limbs.

B. malayi is a mosquito-borne filarial worm. Mosquitos deposit infective third stage larvae (L3) on human skin. The larvae then penetrate and migrate to the lymphatic vessels where they develop into adult worms over several months. Development includes molting transitions into fourth stage larvae (L4) and juvenile adults to reach maturation. The matured female worms release large numbers of microfilariae into the host bloodstream. The microfilariae are ingested by a mosquito during a blood meal and penetrate the midgut and develop over a period of 10 to 14 days to L3. L3 are developmentally arrested in the mosquito. The process repeats when the mosquito's proboscis penetrates human skin.⁴

Material Provided:

NR-48889 consists of up to 2 million *B. malayi* microfilariae. If more material is required for your intended use, please contact BEI Customer Services at <u>contact@beiresources.org</u>, to request the additional material.

Packaging/Storage:

NR-48889 was packaged in 1.5 mL centrifuge tubes. The product is provided on dry ice and should be stored at -20°C or colder immediately upon arrival.

Citation:

Acknowledgment for publications should read "The following reagent was provided by the NIH/NIAID Filariasis Research Reagent Resource Center for distribution by BEI Resources, NIAID, NIH: *Brugia malayi* Microfilariae from the Peritoneal Cavity of a Mongolian Gerbil (Frozen), NR-48889."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. <u>Biosafety in</u> <u>Microbiological and Biomedical Laboratories</u>. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmbl5/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at <u>www.beiresources.org</u>.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC[®] nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC[®] nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC[®] and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC[®], their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, noncommercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a

E-mail: <u>contact@beiresources.org</u> Tel: 800-359-7370 Fax: 703-365-2898 **b**|**e**|**i** resources

SUPPORTING INFECTIOUS DISEASE RESEARCH

license is required. U.S. Government contractors may need a license before first commercial sale.

References:

- Ash, L. R. and J. M. Riley. "Development of Subperiodic Brugia malayi in the Jird, Meriones Unguiculatus, with Notes on Infections in Other Rodents." <u>J. Parasitol.</u> 56 (1970): 969-973. PubMed: 5504534.
- Michalski, M. L., et al. "The NIH-NIAID Filariasis Research Reagent Resource Center." <u>PLoS Negl. Trop.</u> <u>Dis.</u> 5 (2011): e1261. PubMed: 22140585.
- Simonsen, P. E. and M. E. Mwakitalu. "Urban Lymphatic Filariasis." <u>Parasitol. Res.</u> 112 (2013): 35-44. PubMed: 23239094.
- Li, B. W., et al. "Transcription Profiling Reveals Stageand Function-Dependent Expression Patterns in the Filarial Nematode *Brugia malayi*." <u>BMC Genomics</u> 13 (2012): 184. PubMed: 22583769.

ATCC[®] is a trademark of the American Type Culture Collection.

