biei resources

SUPPORTING INFECTIOUS DISEASE RESEARCH

Guinea Pig Expression Clone CD8α/pcDNA3.1 Hygro(+)

Catalog No. NR-36204

For research use only. Not for human use.

Contributor:

Roger G. Rank, Ph.D., Professor, Microbiology and Immunology, College of Medicine, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA

Manufacturer:

BEI Resources

Product Description:

NR-36204 is a full-length expression clone containing the CD8 α gene (GenBank: <u>NM 001172876</u>) from guinea pig (*Cavia porcellus*). The CD8 α gene was cloned into vector pcDNA3.1/hygro(+) via *Bam*HI and *Eco*RV insertion sites, transformed into MAX Efficiency[®] *Escherichia coli* (*E. coli*) DH5 α TM competent cells (InvitrogenTM), and extracted using a QIAGEN[®] Plasmid Mega Kit.

<u>Note</u>: Sequencing results for NR-36204 revealed that most regions of the vector match the depositor's information, including the ampicillin resistance gene, the hygromycin resistance gene under control of the SV40 promoter, polyadenylation signals, and the pUC and f1 origins of replication. However, the native cytomegalovirus (CMV) promoter has been replaced by a ubiquitin (UbC) promoter upstream of CD8 α ; this replacement also resulted in the loss of the T7 promoter. The resulting size of the plasmid is 6.735 kilobases. The plasmid map and the complete plasmid sequence are provided on the Certificate of Analysis for NR-36204.

CD8 is expressed on cytotoxic T cells and functions as a coreceptor for recognition of major histocompatibility complex class I peptide complexes by the T-cell receptor. The CD8 molecule consists of two subunits (α and β) and exists either as a heterodimer ($\alpha\beta$) or a homodimer ($\alpha\alpha$). The sequence homology of guinea pig CD8 indicates greater homology to human, canine, and feline counterparts than to rodent CD8. As the guinea pig serves as an ideal nonprimate animal model to several human diseases, such as syphilis, tuberculosis, and chlamydial genital and ocular infections, CD8 provides a unique molecular tool for studying cell-mediated immune response.¹

Material Provided:

Each vial of NR-36204 contains approximately 1 μ g of plasmid DNA in TE buffer. The concentration is shown on the Certificate of Analysis.

Note: The vial labels for NR-36204 incorrectly state that the plasmid DNA is provided as an *E. coli* glycerol stock.

NR-36204 is purified plasmid DNA that was extracted from bacteria prior to vialing.

Packaging/Storage:

NR-36204 was packaged aseptically in plastic cryovials. The product is provided frozen on dry ice and should be stored at -20°C or colder immediately upon arrival. Freeze-thaw cycles should be minimized.

Post-Transformation Growth Conditions:

Media:

NR-36204 contains the gene required for ampicillin (Amp) resistance. The recommended concentration of Amp in culture is 100 µg/mL. Luria Bertani (LB) broth or equivalent LB agar or equivalent <u>Incubation</u>: Temperature: 37°C Atmosphere: Aerobic <u>Propagation</u>:

Incubate the tube, slant and/or plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Guinea Pig Expression Clone CD8α/pcDNA3.1 Hygro(+), NR-36204."

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. <u>Biosafety in</u> <u>Microbiological and Biomedical Laboratories</u>. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see <u>www.cdc.gov/biosafety/publications/bmbl5/index.htm</u>.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at <u>www.beiresources.org</u>.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC[®] nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC[®] nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. $ATCC^{\circledast}$ and the U.S. Government are not liable for any

E-mail: <u>contact@beiresources.org</u> Tel: 800-359-7370 Fax: 703-365-2898 biei resources

SUPPORTING INFECTIOUS DISEASE RESEARCH

damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC[®], their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, noncommercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

 Nagarajan, U. M., C. O'Connell, and R. G. Rank. "Molecular Characterization of Guinea-Pig (*Cavia porcellus*) CD8α and CD8β cDNA." <u>Tissue Antigens</u> 63 (2004): 184-189. PubMed: 14705990.

ATCC[®] is a trademark of the American Type Culture Collection.

