

Certificate of Analysis for NR-46198

Staphylococcus aureus, Strain CO-84

Catalog No. NR-46198

Product Description: Staphylococcus aureus (S. aureus), strain CO-84 was isolated in 2005 from the blood of an 84-year-old female with a bloodstream infection (BSI) in Colorado, USA. S. aureus, strain CO-84 is a clinically-associated methicillin-resistant S. aureus (MRSA) strain.

Lot¹: 64044896 Manufacturing Date: 11FEB2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology	Gram-positive cocci	Gram-positive cocci
Colony morphology ²	Report results	Circular, low convex, entire, smooth and white (Figure 1)
Motility (wet mount)	Report results	Non-motile
Hemolysis ²	Report results	β-hemolytic
Biochemical characterization	·	
Catalase	Positive	Positive
Coagulase ³	Report results	Positive
VITEK® 2 Compact (GP card)	≥ 90% probability of being S. aureus	S. aureus (99% probability)4
VITEK® MS (MALDI-TOF)	S. aureus	S. aureus (99.9%)
Antibiotic Susceptibility Profile		
VITEK® (AST-GP71 card)⁵		
Beta-lactamase ⁶	Report results	Positive
Cefoxitin screen	Report results	Positive
Benzylpenicillin	Report results	Resistant (≥ 0.5 μg/mL)
Oxacillin	Resistant	Resistant (≥ 4 μg/mL)
Gentamicin	Sensitive	Sensitive (≤ 0.5 μg/mL)
Ciprofloxacin	Report results	Resistant (≥ 8 μg/mL)
Levofloxacin	Resistant	Resistant (≥ 8 µg/mL)
Moxifloxacin	Report results	Resistant (= 4 μg/mL)
Clindamycin (inducible resistance)	Report results	Negative
Erythromycin	Resistant	Resistant (≥ 8 μg/mL)
Clindamycin	Resistant	Resistant (≥ 8 µg/mL)
Quinupristin/dalfopristin	Report results	Sensitive (= 0.5 µg/mL)
Linezolid	Sensitive	Sensitive (= 2 µg/mL)
Daptomycin	Sensitive	Sensitive (= 0.5 µg/mL)
Vancomycin	Sensitive	Sensitive (= 1 µg/mL)
Minocycline	Report results	Sensitive (≤ 0.5 μg/mL)
Tetracycline	Sensitive	Sensitive (≤ 1 μg/mL)
Tigecycline	Report results	Sensitive (≤ 0.12 µg/mL) ⁷
Nitrofurantoin	Report results	Sensitive (≤ 16 μg/mL)
Rifampicin	Sensitive	Sensitive (≤ 0.5 µg/mL)
Trimethoprim/sulfamethoxazole	Sensitive	Sensitive (≤ 10 μg/mL)
Etest® antibiotic test strips8		
Chloramphenicol ⁹	Report results	Sensitive (= 6 µg/mL) ¹⁰
Teicoplanin ⁹	Report results	Sensitive (= 1.5 μg/mL)
Genotypic Analysis		
Sequencing of 16S ribosomal RNA gene	≥ 99% sequence identity to S. aureus	100% sequence identity to S. aureus
(~ 1500 base pairs)	type strain (GenBank: L37597)	type strain (GenBank: L37597)
Purity (post-freeze) ¹¹	Consistent with expected colony morphology	Consistent with expected colony morphology
Viability (post-freeze) ²	Growth	Growth

BEI Resources www.beiresources.org E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898



SUPPORTING INFECTIOUS DISEASE RESEARCH

Certificate of Analysis for NR-46198

²1 day at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood

³4 hours at 37°C in rabbit serum with 0.15% EDTA (Coagulase Plasma BBL™ 240827)

¹¹Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood.





Date: 21 JUN 2016 Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

¹S. aureus, strain CO-84 was deposited to BEI Resources as part of the NARSA collection. NR-46198 was produced by inoculation of the deposited material into Tryptic Soy broth and grown 1 day at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar with 5% defibrinated sheep blood kolles which were grown 1 day at 37°C in an aerobic atmosphere to produce this lot.

⁴Percent probabilities above 90% indicate a close match to the typical biochemical pattern for the given organism, with a percent probability of 99% being a perfect match between the test reaction pattern and the unique biochemical pattern of the given organism or organism group. For additional information, please refer to O'Hara, C.M. and J. M. Miller. "Evaluation of the Vitek 2 ID-GNB Assay for Identification of Members of the Family Enterobacteriaceae and Other Nonenteric Gram-Negative Bacilli and Comparison with the Vitek GNI+ Card." J. Clin. Microbiol. 41 (2003): 2096-2101. PubMed: 12734254.

⁵Minimum Inhibitory Concentration (MIC); MIC Interpretation Guideline: CLSI M100-S22 (2012)

⁶The production of beta-lactamase was detected using a Cefinase™ Paper Disc (BBL™ 231650).

⁷MIC Interpretation Guideline: EUCAST Version 4.0 (2014)

⁸¹ day at 37°C in an aerobic atmosphere on Mueller Hinton agar

⁹For both chloramphenicol (bioMérieux Etest® 412308) and teicoplanin (bioMérieux Etest® 412459), a MIC ≤ 8 μg/mL is sensitive, a MIC = 16 μg/mL is intermediate and a MIC ≥ 32 μg/mL is resistant.

¹⁰S. aureus, strain CO-84 was deposited as having an intermediate susceptibility to chloramphenicol. Antibiotic susceptibility testing performed in duplicate determined that strain CO-84 is susceptible to chloramphenicol.