

Certificate of Analysis for NR-43661

Helicobacter pylori, Strain Hp H-27

Catalog No. NR-43661

Product Description: *Helicobacter pylori* (*H. pylori*), strain Hp H-27 was isolated from gastric biopsy homogenate of a patient with a gastric ulcer in Ohio, USA.

Lot¹: 64136558 Manufacturing Date: 06APR2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology	Gram-negative rods	Gram-negative rods
Colony morphology ²	Report results	Circular, low convex, entire, translucent and gray (Figure 1)
Motility (wet mount)	Report results	Motile gray (Figure 1)
Biochemical characterization	Positive	Desitive
Catalase Oxidase	Positive	Positive Positive
		Positive
Urease	Positive	
Nitrate reduction	Negative	Negative
H2S (lead acetate paper)	Report results	Positive
Hippurate hydrolysis	Negative	Negative
Growth with 5% CO ₂	Growth	Growth
Growth at 25°C	No growth	No growth
Growth at 37°C	Growth	Growth
Growth at 42°C	Report results	Growth
Brucella albimi + 0.16% agar (growth control)	Growth	Growth
Brucella albimi + 0.16% agar with 1% glycine	No growth	Growth ³
Brucella albimi + 0.16% agar with 3.5% NaCl	No growth	Growth ⁴
Antibiotic Susceptibility Profile		
BD BBL™ Sensi-Disc™ susceptibility test discs		
Metronidazole (80 μg) ⁵	Report results	50 mm
Nalidixic acid (30 μg) ⁶	Report results	8 mm
Genotypic Analysis	≥ 99% sequence identity to	99.9% sequence identity to
Sequencing of 16S ribosomal RNA gene	<i>H. pylori,</i> strain Hp H-27	H. pylori, strain Hp H-27
(~ 710 base pairs)	(GenBank: AKOH01000001.1)	(GenBank: AKOH01000001.1)
Confirmation of <i>H. pylori</i> by PCR Amplification of	,	
Extracted DNA		
Positive control (16S ribosomal RNA gene)	Amplicon present	Amplicon present
Negative control (H. acinonychis)	No amplicon present	No amplicon present
ureA	Amplicon present	Amplicon present
ssaA	Amplicon present	Amplicon present
	Amphoon present	Amplicon present
Purity (post-freeze)		
Microaerophilic growth ⁷	Growth consistent with expected	Growth consistent with expected
	colony morphology	colony morphology
Aerobic growth ^{8,9}	Growth consistent with expected	Growth consistent with expected
	colony morphology	colony morphology
Viability (post-freeze) ²	Growth	Growth

¹NR-43661 was produced by inoculation of the deposited material into Brucella broth. Broth inoculum was added to Columbia agar with 7% defibrinated horse blood, 5 μg/mL trimethoprim, 5 μg/mL vancomycin, 10 μg/mL cefsulodin and 2.5 μg/mL amphotericin B. The inoculated agar and broth were each grown for 3 days at 37°C in a microaerophilic atmosphere (~ 6-16% O₂ and 2-10% CO₂). Colonies from the Columbia agar culture were suspended into the Brucella broth growth, and this biphasic culture was added to Columbia agar with 7% defibrinated horse blood, 5 μg/mL trimethoprim, 5 μg/mL vancomycin, 10 μg/mL cefsulodin and 2.5 μg/mL amphotericin B kolles, which were grown for 2 days 37°C in a microaerophilic atmosphere to produce this lot.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898



Certificate of Analysis for NR-43661

²2 days on Columbia agar with 7% defibrinated horse blood, 5 μg/mL trimethoprim, 5 μg/mL vancomycin, 10 μg/mL cefsulodin and 2.5 μg/mL amphotericin B at 37°C in a microaerophilic atmosphere

³Specifications for these tests were obtained from Bergey's Manual[®] of Systematic Bacteriology, 2nd ed., Volume 2, Part C, which indicates that growth may occur in up to 17% of strains.

⁴Specifications for these tests were obtained from Bergey's Manual® of Systematic Bacteriology, 2nd ed., Volume 2, Part C, which indicates that growth may occur in 20% to 43% of strains.

⁵Test performed using metronidazole 80 µg (MET-80, BBL™ catalog no. 231605)

⁶Test performed using nalidixic acid 30 µg (NA-30, BBL™ catalog no. 231311)

⁷Purity of this lot was assessed for 7 days on Columbia agar with 7% defibrinated horse blood, 5 μg/mL trimethoprim, 5 μg/mL vancomycin, 10 μg/mL cefsulodin and 2.5 μg/mL amphotericin B at 37°C in a microaerophilic atmosphere (~ 6-16% O₂ and 2-10% CO₂).

⁸Purity of this lot was assessed for 7 days on Tryptic Soy agar with 5% defibrinated sheep blood at 37°C in an aerobic atmosphere with 5% CO₂.

⁹H. pylori is known to show weak growth under aerobic conditions (Bury-Moné, S., et al. "Is *Helicobacter pylori* a True Microaerophile?" <u>Helicobacter</u> 11 (2006): 296-303. PubMed: 16882333.).

Figure 1: Colony Morphology

Date: 20 JUL 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898