

Escherichia coli, Strain SS724

Catalog No. NR-50519

For research use only. Not for human use.

Contributor:

James B. Kaper, Associate Dean, Department of Microbiology and Immunology, University of Maryland, Baltimore, Maryland, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: Enterobacteriaceae, Escherichia

Species: Escherichia coli

Strain: SS724

Serotype: O127:H6

Original Source: Escherichia coli (E. coli), strain SS724 is a glutamase decarboxylase (gadX) insertion mutant of the wild type strain E2348/69.¹ Strain E2348/69 was isolated in 1969 during an outbreak of diarrhea in an infant nursery in Taunton, England.²⁻⁴

Comments: Mutagenesis occurred through the insertion of a kanamycin-resistance gene (aphT) cassette from pUC18K into gadX.¹ gadX encodes for a transcription regulator, GadX, which may be involved in regulation of genes required for acid resistance and virulence including per, gadA and gadB.¹

E. coli is a Gram-negative, rod-shaped bacterium commonly found in the gut flora of warm-blooded animals and is the primary facultative anaerobe of the human gastrointestinal tract. There are a number of pathogenic types of E. coli associated with diarrhea that are referred to as: enterohemorrhagic E. coli (EHEC) [also known as Shiga toxin-producing E. coli (STEC) or Verocytotoxin-producing E. coli (VTEC)]⁵, enterotoxigenic E. coli (ETEC)⁶, enteropathogenic E. coli (EPEC)⁷, enteroaggregative E. coli (EAEC)⁸, enteroinvasive E. coli (EIEC) and diffusely adherent E. coli (DAEC).⁹

Characteristic features of EPEC strains are induction of attaching and effacing (A/E) lesions on intestinal epithelial cells, lack of enterotoxins and lack of shigella-like invasiveness. The ability to induce A/E lesions is encoded by genes located on a 35-kb pathogenicity island (PAI) called the locus of enterocyte effacement (LEE), which contains the genes encoding eae (intimin), a type III secretion system, a number of secreted proteins (ESP), and the translocated intimin receptor (Tir).⁷

EPEC strain E2348/69 (serotype O127:H6) has been used worldwide as a prototype strain to study EPEC biology, genetics, and virulence. The complete genome sequence of strain E2348/69 (GenBank: [NC_011601](#)) has enabled analysis of over 400 known/predicted effector sequences and identified only 21 putative effectors, providing a clear picture of the core LEE and non-LEE effector genes.³

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Tryptic Soy broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-50519 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy broth or Nutrient broth or equivalent

Tryptic Soy agar or Nutrient agar or Tryptic Soy agar with 5% defibrinated sheep blood or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate at 37°C for 1 day.

Citation:

Acknowledgment for publications should read "The following reagent was provided by Dr. Kaper, for distribution by BEI Resources, NIAID, NIH: Escherichia coli, Strain SS724, NR-50519."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370

Fax: 703-365-2898

Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Shin, S., et al. "An Activator of Glutamate Decarboxylase Genes Regulates the Expression of Enteropathogenic *Escherichia coli* Virulence Genes through Control of the Plasmid-Encoded Regulator, Per." *Mol. Microbiol.* 41 (2001): 1133-1150. PubMed: 11555293.
2. Levine, M. M., et al. "The Diarrheal Response of Humans to Some Classic Serotypes of Enteropathogenic *Escherichia coli* is Dependent on a Plasmid Encoding an Enteroadhesiveness Factor" *J. Infect. Dis.* 152 (1985): 550-559. PubMed: 2863318.
3. Iguchi, A., et al. "Complete Genome Sequence and Comparative Genome Analysis of Enteropathogenic *Escherichia coli* O127:H6 Strain E2348/69." *J. Bacteriol.* 191 (2009): 347-354. PubMed: 18952797.
4. Nataro, J. P. and J. B. Kaper. "Diarrheagenic *Escherichia coli*." *Clin. Microbiol. Rev.* 11 (1998): 142-201. PubMed: 9457432.
5. Smith, J. L., P. M. Fratamico and N. W. Gunther IV. "Shiga Toxin-Producing *Escherichia coli*." *Adv. Appl. Microbiol.* 86 (2014): 145-197. PubMed: 24377855.
6. Zhang, W. and D. A. Sack. "Progress and Hurdles in the Development of Vaccines against Enterotoxigenic *Escherichia coli* in Humans." *Expert Rev. Vaccines* 11 (2012): 677-694. PubMed: 22873126.
7. Ochoa, T. J. and C. A. Contreras. "Enteropathogenic *Escherichia coli* Infection in Children." *Curr. Opin. Infect. Dis.* 24 (2011): 478-483. PubMed: 21857511.
8. Estrada-Garcia, T. and F. Navarro-Garcia. "Enterotoxigenic *Escherichia coli* Pathotype: A Genetically Heterogeneous Emerging Foodborne Enteropathogen." *FEMS Immunol. Med. Microbiol.* 66 (2012): 281-298. PubMed: 22775224.
9. Smith, E. J., et al. "Pathogenesis of Adherent-Invasive *Escherichia coli*." *Future Microbiol.* 8 (2013): 1289-1300. PubMed: 24059919.

ATCC® is a trademark of the American Type Culture Collection.

