

**Alphacoronavirus 1, Purdue P115 (attenuated)
(formerly Porcine Transmissible Gastroenteritis Virus)**

Catalog No. NR-43285

Derived from BEI Resources NR-446

Product Description: Clarified cell lysate and supernatant from *Sus scrofa* testicular fibroblasts¹ infected with alphacoronavirus 1, Purdue P115 (attenuated)

Lot²: 70007016

Manufacturing Date: 14JUL2017

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in ST Cells ¹	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Species-Specific Region (406 nucleotides)	Consistent with alphacoronavirus 1, Purdue P115	99% identity with alphacoronavirus 1, Purdue P115 (GenBank: DQ811788)
Titer by TCID ₅₀ Assay ^{3,4} in ST Cells ¹	Report results	2.8 × 10 ⁷ TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

¹ST cells (ATCC® CRL-1746™)

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 2 days at 37°C and 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴5 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

13 FEB 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

