

## *Yersinia pestis*, Strain Yokohama Derivative (D10)

**Catalog No. NR-4692**

**Product Description:** *Yersinia pestis* (*Y. pestis*) is an aerobic, non-spore-forming, Gram-negative rod-shaped bacterium. *Y. pestis*, strain Yokohama D10 is a derivative of strain Yokohama which contains the pMT1 and the pPCP1 plasmids, and the unstable chromosomal *pgm* locus but lacks the pCD1 plasmid that is essential for virulence.

**Lot<sup>1</sup>: 57753203**

**Manufacturing Date: 20JUL2007**

TEST	SPECIFICATIONS	RESULTS
<b>Phenotypic Analysis</b> Cellular morphology Colony morphology <sup>2</sup>  Congo red (CR) agar <sup>3,4</sup> Biochemical Analyses Analytical profile index (API 20 E®) Nitrate reduction Fermentation of glycerol Urease	Gram-negative rod Report results  Red colonies (Crb <sup>+</sup> )  Consistent with <i>Y. pestis</i> Positive Positive Negative	Gram-negative rod Circular, entire, convex, opaque (Figure 1) Red colonies (Crb <sup>+</sup> )  Consistent with <i>Y. pestis</i> Positive Positive Negative
<b>Genotypic Analysis</b> Sequencing of 16S ribosomal RNA gene (~ 1420 bp)	Consistent with <i>Y. pestis</i>	Consistent with <i>Y. pestis</i> <sup>5</sup>
<b>PCR Assay of Extracted DNA</b> 16S ribosomal RNA gene Presence of virulence plasmids pMT1 (~ 100 kb plasmid) pCD1 (~ 70 kb plasmid) pPCP1 (~ 9.5 kb plasmid)	~ 1500 bp amplicon  ~ 1200 bp amplicon Not present in this derivative ~ 400 bp amplicon	~ 1500 bp amplicon  ~ 1200 bp amplicon No product amplified ~ 400 bp amplicon
<b>Viability (post-freeze)<sup>2</sup></b>	Growth on agar	Growth on agar

<sup>1</sup>*Y. pestis*, strain Yokohama (D10) was deposited by Professor Robert R. Brubaker of the Department of Microbiology and Molecular Genetics at Michigan State University, East Lansing, Michigan. NR-4688 was prepared by broth (Tryptic Soy Broth; BD 211768) culture of the deposited material and grown 48 hours at 28°C and aerobic atmosphere. Broth inoculum was added to Kolles which were grown 48 hours at 28°C and aerobic atmosphere to produce this lot.

<sup>2</sup>48 hours at 28°C and aerobic atmosphere on Tryptic Soy Agar (BD 236950)

<sup>3</sup>1 to 4 days at 28°C and aerobic atmosphere on CR agar

<sup>4</sup>Hare, J. M. and K. A. McDonough. "High-Frequency RecA-Dependent and -Independent Mechanisms of Congo Red Binding Mutations in *Yersinia pestis*." *J. Bacteriol.* 181 (1999): 4896-4904. PubMed: 10438760.

<sup>5</sup>Also consistent with other *Yersinia* species

**Figure 1**



**Date:** 07 AUG 2008

**Signature:** Signature on File

**Title:** Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.  
You are authorized to use this product for research use only. It is not intended for human use.

