

Chikungunya Virus, H 20235-St. Martin-2013

Catalog No. NR-49901

Product Description: Chikungunya virus (CHIKV), H 20235-St. Martin-2013 was isolated from a human in Saint Martin in 2013. Each vial contains cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells¹ infected with CHIKV, H 20235-St. Martin-2013.

Passage History: XV2/V3 (Prior to deposit at BEI Resources/BEI Resources); X = Unknown; V = Vero cells

Lot²: 70013683

Manufacturing Date: 16MAR2018

TEST	SPECIFICATIONS	RESULTS
Infectivity in Vero E6 Cells¹	Report results	Cell rounding and detachment
Sequencing of Species-Specific Region (~ 930 nucleotides)	Consistent with CHIKV, H 20235-St. Martin-2013	100% identity with CHIKV, H 20235-St. Martin-2013 (GenBank: MG208125)
Titer by TCID₅₀ Assay^{3,4} in Vero E6 Cells¹ by Cytopathic Effect	Report results	2.8 × 10 ⁶ TCID ₅₀ per mL
Amplification of CHIKV Sequence by RT-PCR	~ 1070 base pair amplicon	~ 1070 base pair amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Vero 76, clone E6; ATCC® CRL-1586™

²Grown in Eagle's Minimum Essential Medium containing Earle's Balanced Salt Solution, non-essential amino acids, 2 mM L-glutamine, 1 mM sodium pyruvate and 1.5 g/L of sodium bicarbonate (ATCC® 30-2003) supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 3 days at 37°C with 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴5 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/
Heather Couch

06 AUG 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.



BEI Resources
www.beiresources.org

E-mail: contact@beiresources.org
Tel: 800-359-7370
Fax: 703-365-2898