

Vector pMCSG53 Containing the SARS-Related Coronavirus 2, Wuhan-Hu-1 Non-Structural Protein 15 Gene

Catalog No. NR-52426

This reagent is the tangible property of the U.S. Government.

Product Description:

The non-structural protein 15 (nsp15) gene from severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), Wuhan-Hu-1 (GenBank: [MN908947](#)) was codon optimized and cloned into the pMCSG53 plasmid. pMCSG53 is an *Escherichia coli* (*E. coli*) expression vector that contains an N-terminal hexa-histidine tag, followed by a tobacco etch virus (TEV) protease recognition site prior to the insert coding sequence, resulting in the expression of a cleavable histidine-tagged protein. It also contains tRNA genes covering rare codons for Arg (AGG/AGA) and Ile (AUA) to improve expression in *E. coli*. The beta-lactamase gene, *bla*, provides transformant selection through ampicillin resistance in *E. coli*. The deposited plasmid was transformed into One Shot™ TOP10 *Escherichia coli* (Invitrogen™ C404003), grown in Luria-Bertani broth with ampicillin (50 µg per mL) for 1 day at 37°C in an aerobic atmosphere, extracted using a Plasmid Plus Maxi Kit (QIAGEN® 12963) and vialled in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0).

Lot: 70035121

Manufacturing Date: 06MAY2020

| TEST | SPECIFICATIONS | RESULTS |
|--|---|--|
| Next-Generation DNA Sequencing | ~ 5850 base pairs | 5849 base pairs ¹ |
| Genotypic Analysis Sequencing of nsp15 insert (~ 1040 base pairs) N-terminal His ₆ tag N-terminal TEV protease site | 100% sequence identity to depositor's sequence His ₆ tag sequence confirmed TEV protease site sequence confirmed | 100% sequence identity to depositor's sequence ² His ₆ tag sequence confirmed TEV protease site sequence confirmed |
| Antibiotic Resistance Ampicillin (encoded by beta-lactamase gene <i>bla</i>) ³ | <i>bla</i> sequence present | <i>bla</i> sequence present |
| Concentration by PicoGreen® Measurement | ≥ 2 µg/mL | 0.2 µg in 20 µL per vial (12 µg/mL) |
| Amount per Vial | Report results | 0.2 µg per vial |
| OD₂₆₀/OD₂₈₀ Ratio (pre-vial) | 1.7 to 2.1 | 1.9 |
| Effective Bacterial Transformation Invitrogen™ One Shot™ TOP10 <i>Escherichia coli</i> | ≥ 50 colonies per ng | 68 colonies per ng |

¹The sequence was assembled pre-vial using the depositor's predicted sequence as the reference sequence. The complete plasmid sequence and map are provided on the BEI Resources webpage.

²The NR-52426 insert was codon optimized, but otherwise is consistent with the SARS-CoV-2, Wuhan-Hu-1 nsp15 protein (YP_009725310.1).

³The antibiotic ampicillin degrades quickly during growth. Bacterial stationary phase should be minimized during plasmid expansion to avoid plasmid loss and increased antibiotic concentrations may be necessary.

/Heather Couch/

Heather Couch

28 MAY 2020

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by ATCC® and the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

