



NIH AIDS Reagent Program

20301 Century Boulevard
Building 6, Suite 200
Germantown, MD 20874
USA

Phone: 240 686 4740
Fax: 301 515 4015
aidsreagent.org

DATA SHEET

Reagent:	Een217 T Cell Clone
Catalog Number:	755
Lot Number:	3/19/90
Release Category:	A
Provided:	5 x 10 ⁶ cells/vial.
Propagation Medium:	RPMI 1640 supplemented with 10% heat-inactivated fetal bovine serum, 4 mM L-glutamine, 50 U/mL penicillin, 50 µg/mL streptomycin, 50 U/mL recombinant human IL-2
Freeze Medium:	Fetal calf serum, 90%; DMSO, 10%.
Growth Characteristics:	For continued growth, these cells must be re-stimulated every 7-14 days with PHA and irradiated allogeneic peripheral blood mononuclear cells. Please see cell propagation instructions attached to data sheet.
Sterility:	Negative for aerobic and anaerobic bacteria, mycoplasma, fungi, and yeast.
Description:	CD4+ gp120-specific human cytolytic T-cell clone.
Special Characteristics:	This clone was derived from an HIV-seronegative donor by <i>in vitro</i> stimulation with recombinant gp120 followed by soft agar cloning. It recognizes aa 410-429 of HIV-1 pV22 gp120 in association with certain subtypes of DR4 (Dw10 and Dw15), and is cytolytic. Cells are 99% CD4+, 99% CD3+.
	<u>PROPAGATION OF HUMAN T-CELL CLONES</u>
Recommended Storage:	Liquid nitrogen.

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

Contributor: Dr. Robert F. Siliciano.

References: Preparation of irradiated PBMCs: Enhanced culture assay for detection and quantitation of latently infected, resting CD4+ T-cells carrying replication-competent virus in HIV-1-infected individuals. Siliciano JD, Siliciano RF. *Methods Mol Biol.* 2005;304:3-15.[Abstract](#)

Siliciano RF, Lawton T, Knal C, Karr R, Berman P, Gregory T, Reinherz E. Analysis of host-virus interactions in AIDS with anti-gp120 T cell clones: Effects of HIV sequence variation and a mechanism for CD4+ cell depletion. *Cell* **54**:561-575, 1988.

Callahan K, Fort M, Obah E, Reinherz E, Siliciano, R. Genetic variability in HIV-1 gp120 affects interactions with HLA molecules and T cell receptor. *J Immunol* **144**:3341-3346, 1990.

Polydefkis M, Koenig S, Flexner C, Obah E, Gebo K, Chakrabarti S, Earl P, Moss B, Siliciano R. Anchor sequence-dependent endogenous processing of human immunodeficiency virus 1 envelope glycoprotein gp160 for CD4+ T cell recognition. *J Exp Med* **171**:875-887, 1990.

NOTE: Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Een217 T Cell Clone from Dr. Robert F. Siliciano." Also include the references cited above in any publications.

Last Updated July 02, 2018

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.