

**Influenza B Virus, B/Brisbane/33/2008 (Victoria Lineage)**

**Catalog No. NR-42006**

Derived from CDC ID No. 2009715789

**Product Description:**

Influenza B virus, B/Brisbane/33/2008 (Victoria Lineage) was isolated from a human in Brisbane, Australia in 2008. NR-42006 lot 70041362 is derived from CDC ID No. 2009715789 and was produced in the allantoic cavity of specific pathogen free (SPF) embryonated chicken eggs (9- to 11-day-old; Charles River, Norwich, Connecticut, USA) infected with seed material for 2 days at 34°C in a humidified chamber.

**Passage History:**

E(3)/E(4)/E(1)/E(2) (Submission Laboratory/Centers for Disease Control and Prevention/International Reagent Resource/BEI Resources); E = SPF embryonated chicken eggs

**Lot: 70041362**

**Manufacturing Date: 14JAN2021**

TEST	SPECIFICATIONS	RESULTS
<b>Identification by Infectivity Using Embryonated Chicken Eggs</b> Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
<b>Sequencing of Hemagglutinin and Matrix Coding Regions</b> Hemagglutinin (~ 930 nucleotides)	≥ 98% identity with B/Brisbane/33/2008 (Victoria Lineage) (GenBank: CY149981.1)	99.9% identity with B/Brisbane/33/2008 (Victoria Lineage) (GenBank: CY149981.1)
<b>Titer by CEID<sub>50</sub> Assay in Embryonated Chicken Eggs<sup>1</sup></b> (2 days at 34°C in a humidified chamber)	Report results	1.6 × 10 <sup>7</sup> CEID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>2</sup> Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>The Chicken Embryo Infectious Dose 50% (CEID<sub>50</sub>) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID<sub>50</sub> provides a measure of the infectious titer (or infectivity) of a virus preparation.

<sup>2</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Heather Couch/

Heather Couch

22 JUL 2021

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

