

Influenza A Virus, A/Uruguay/716/2007 (H3N2)

Catalog No. NR-42003

Derived from CDC ID No. 2007731384

Product Description:

Influenza A virus, A/Uruguay/716/2007 (H3N2) was isolated from a human in Uruguay on June 21, 2007. NR-42003 lot 70051832 was produced in the allantoic cavity of specific pathogen free (SPF) embryonated chicken eggs (10- to 11-day-old; Charles River, Norwich, Connecticut, USA) infected with seed material (BEI Resources lot 61361429) for 2 days at 35°C in a humidified chamber.

Lot: 70051832

Manufacturing Date: 28APR2022

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin and Matrix Coding Regions Hemagglutinin (~ 730 nucleotides) Matrix (~ 950 nucleotides)	≥ 98% identity with A/Uruguay/716/2007 (H3N2) (GenBank: CY121632) ≥ 98% identity with A/Uruguay/716/2007 (H3N2) (GenBank: EU716434)	99.9% identity with A/Uruguay/716/2007 (H3N2) (GenBank: CY121632) 100% identity with A/Uruguay/716/2007 (H3N2) (GenBank: EU716434)
Titer by CEID₅₀ Assay in Embryonated Chicken Eggs¹ (2 days at 35°C in a humidified chamber)	Report results	8.9 × 10 ⁷ CEID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ² Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

²Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorum Brower/
Sonia Bjorum Brower

29 JUN 2022

Lead Technical Writer or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

