

# **Certificate of Analysis for NR-56773**

# Rhinovirus A60, 2268-CV37

# Catalog No. NR-56773

# **Product Description:**

Rhinovirus A60, 2268-CV37 was isolated prior to 1963. NR-56773 lot 70053414 was produced by infecting *Homo sapiens* lung fibroblasts (WI-38; ATCC® CCL-75™) with Rhinovirus A60, 2268-CV37 and incubating in Eagle's Minimum Essential Medium (EMEM; ATCC® 30-2003™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 8 days at 33°C with 5% CO₂.

# Passage History:

H(5)WI(5)H(2)/WI(1) (Prior to deposit/BEI Resources); H = HeLa; WI = WI-38 cells

Lot: 70053414 Manufacturing Date: 01JUL2022

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in WI-38 Cells	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Species-Specific Region (~ 780 nucleotides)	≥ 98% identity with HRV A60, 2268-CV37 (GenBank: FJ445143.1)	100% identity with HRV A60, 2268-CV37 (GenBank: FJ445143.1)
Titer by TCID <sub>50</sub> Assay in WI-38 Cells by Cytopathic Effect and Immunofluorescence <sup>1,2</sup> (8 days at 33°C with 5% CO <sub>2</sub> )	Report results	2.8 × 10 <sup>6</sup> TCID <sub>50</sub> per mL
Sterility (21-day incubation)		
Harpo's HTYE broth, 37°C and 26°C, aerobic <sup>3</sup>	No growth	No growth
Trypticase Soy broth, 37°C and 26°C, aerobic	No growth	No growth
Sabouraud broth, 37°C and 26°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, anaerobic	No growth	No growth
Thioglycollate broth, 37°C, anaerobic	No growth	No growth
DMEM with 10% FBS, 37°C, aerobic	No growth	No growth
Mycoplasma Contamination		
Agar and broth culture (14-day incubation at 37°C)	None detected	None detected
DNA detection by PCR of extracted Test Article nucleic acid	None detected	None detected

<sup>&</sup>lt;sup>1</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation. <sup>2</sup>Pan Enterovirus Reagent (Blend) (Light Diagnostics cat no. 3360)

/Sonia Bjorum Brower/ Sonia Biorum Brower

10 NOV 2022

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Tel: 800-359-7370 Fax: 703-365-2898

<sup>&</sup>lt;sup>3</sup>Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.