

Certificate of Analysis for NR-10462

Acanthamoeba lenticulata, Strain 72/2

Catalog No. NR-10462

Product Description: Acanthamoeba lenticulata (A. lenticulata), strain 72/2 was isolated from the nasal mucosa of a healthy individual in Germany.

Lot¹: 59741149 Manufacturing Date: 10FEB2011

TEST	SPECIFICATIONS	RESULTS
Genotyping Sequencing of 118S ribosomal RNA gene (~ 400 bp)	Consistent with A. lenticulata	Consistent with A. lenticulata
Functional Activity by PCR Amplification ² 18S ribosomal RNA gene (amplicon ASA.S1)	423 bp to 551 bp amplicon	~ 450 bp amplicon
Viable Cell Count by Hemacytometry (pre-freeze)	> 10 ⁶ cells/mL	1.8 x 10 ⁶ cells/mL
Viability (post-freeze) ³	Growth	Growth

The deposited material was cultivated at 20°C and an aerobic atmosphere in fresh water amoeba agar medium (ATCC medium 997) inoculated with Enterobacter aerogenes (ATCC® 13048TM) and preserved. NR-10462 was produced by cultivation of the preserved material under the propagation conditions described above.

Date: 15 JUN 2012 Signature:

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²PCR amplification was performed using the JDP1 and JDP2 primer set as described [Schroeder, J. M. et al. "Use of Subgenic 18S Ribosomal DNA PCR and Sequencing for Genus and Genotype Identification of Acanthamoebae from Humans with Keratitis and from Sewage Sludge." <u>J. Clin. Microbiol.</u> 39 (2001): 1903-1911. PubMed: 11326011].

³Viable cells and signs of infection were seen after 5 days under cultivation conditions at 20°C