

Certificate of Analysis for NR-33661

Acanthamoeba sp., Strain CDC:V155

Catalog No. NR-33661

Product Description: *Acanthamoeba* sp., strain CDC:V155 was isolated in 1989 from the water in a hot spring located in California, USA.

Lot¹: 61178082 Manufacturing Date: 23AUG2012

TEST	SPECIFICATIONS	RESULTS
Genotyping Sequencing of 18S ribosomal RNA gene (~ 440 bp)	Consistent with Acanthamoeba sp.	Consistent with Acanthamoeba sp.
Functional Activity by PCR Amplification ² 18S ribosomal RNA gene (amplicon ASA.S1)	423 bp to 551 bp amplicon	~ 450 bp amplicon
Viable Cell Count by Hemacytometry (pre-freeze)	> 10 ⁶ cells/mL	1.3 x 10 ⁷ cells/mL
Viability (post-freeze) ³	Growth	Growth
Sterility (21-day incubation) Harpo's HTYE broth ⁴ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Brain heart infusion, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic	No growth	No growth

NR-33661 was produced by cultivation of *Acanthamoeba* sp., strain CDC:V155 in PYG Medium (ATCC medium 712) for 8 days at 25°C in an aerobic atmosphere and preserved.

Date: 16 MAY 2013 Signature:

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources
www.beiresources.org

E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²PCR amplification was performed using the JDP1 and JDP2 primer set as described [Schroeder, J. M. et al. "Use of Subgenic 18S Ribosomal DNA PCR and Sequencing for Genus and Genotype Identification of Acanthamoebae from Humans with Keratitis and from Sewage Sludge." <u>J. Clin. Microbiol.</u> 39 (2001): 1903-1911. PubMed: 11326011].

³Viable cells were observed after 1 day under cultivation conditions.

⁴Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.