

Certificate of Analysis for NR-36157

Monoclonal Anti-Hantaan Virus Gc Glycoprotein, Clone 3B6 (produced in vitro)

Catalog No. NR-36157

This reagent is the property of the U.S. Government.

Product Description: Mouse monoclonal antibody prepared against the Hantaan virus Gc (formerly G2) glycoprotein was purified from clone 3B6 hybridoma supernatant by protein G affinity chromatography.

Lot: 61101788 Manufacturing Date: 31AUG2012

TEST	SPECIFICATIONS	RESULTS
Antibody Class Determination	IgG1ĸ	IgG1κ
Experion Pro260 Analysis	Correct molecular weight (MW) for heavy and light chains Report results	Correct MW for heavy and light chains 95.3% pure
Concentration by Spectrophotometer at OD ₂₈₀	Report results	1.1 mg per mL
Functional Activity ¹ Indirect immunofluorescence assay (IFA) titer ² Plaque reduction neutralization test (PRNT ₅₀) titer ³	Report results Report results	256000 ≤ 20
Sterility	0.22 µm filter-sterilized	0.22 µm filter-sterilized

¹Testing performed at U.S. Army Medical Research Institute of Infectious Diseases

Date: 11 FEB 2014 Signature:

Title: Technical Manager, BEI Authentication or designee

Michael Q. Comple

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by ATCC[®] and the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

NR-36157_61101788_11FEB2014

²The IFA titer is defined as the reciprocal of the lowest antibody dilution where glycoprotein staining was clearly visible above background in Hantaan virus 76-118-infected Vero E6 cells.

³The PRNT₅₀ is defined as the reciprocal of the antibody dilution required to reduce the number of Hantaan virus 76-118 plaques in Vero E6 cells by 50% relative to the control wells.