

Certificate of Analysis for NR-46639

Staphylococcus aureus subsp. aureus, Strain JE2, Transposon Mutant NE96 (SAUSA300_2438)

Catalog No. NR-46639

Product Description: Staphylococcus aureus (S. aureus) subsp. aureus, transposon mutant NE96 was derived from S. aureus subsp. aureus, strain JE2. Mutagenesis occurred through the use of the mariner-based transposon bursa aurealis resulting in an erythromycin-resistant deletion strain of JE2. S. aureus subsp. aureus, transposon mutant NE96 was created by disruption of staphylococcal accessory regulator U (sarU), which regulates virulence factors and is a sarA homolog. Strain JE2 is a plasmid-cured derivative of strain LAC that was isolated in 2002 from a skin and soft tissue infection of an inmate in the Los Angeles County Jail in California, USA.

Lot¹: 62489420 Manufacturing Date: 10APR2014

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ² Motility (wet mount)	Gram-positive cocci Report results Report results	Gram-positive cocci Circular, convex, entire, smooth and cream (Figure 1) Non-motile
Confirmation of Transposon Insertion ³	Resistant to erythromycin	Resistant to erythromycin
Viability (post-freeze) ²	Growth	Growth

NR-46639 was produced by inoculation of the deposited material into Tryptic Soy broth with 5 μg/mL erythromycin and incubated for 20 hours at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar with 5 μg/mL erythromycin kolles which were grown 23 hours at 37°C in an aerobic atmosphere to produce this lot.

Figure 1: Colony Morphology



BEI Resources
www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²24 hours at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5 µg/mL erythromycin

³Prior to initiating work, it is recommended that the presence and location of the transposon is confirmed. Gene specific primers should be paired with either the "Upstream" primer (5'-CTCGATTCTATTAACAAGGG-3') for transposons in the "plus" orientation or the "Buster" primer (5'-GCTTTTTCTAAATGTTTTTTAAGTAAATCAAGTAC-3') for transposons in the "minus" orientation. For additional information, refer to Fey, P. D., et al. "A Genetic Resource for Rapid and Comprehensive Phenotype Screening of Nonessential *Staphylococcus aureus* Genes." MBio 4 (2013): e00537-12. PubMed: 23404398.



Certificate of Analysis for NR-46639

Date: 24 SEP 2015

Signature: (

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898