

## **Certificate of Analysis for NR-2595**

## Entamoeba histolytica, HB-301:NIH CL-1-4

Catalog No. NR-2595 (Derived from ATCC® 50548<sup>TM</sup>)

**Product Description:** Entamoeba histolytica is a pathogenic protozoan parasite that predominantly infects humans and other primates. Entamoeba histolytica results in an asymptomatic carrier state in most individuals, but can cause diseases ranging from chronic, mild diarrhea to fulminant dysentery.

Lot<sup>1</sup>: 5078409 Manufacturing Date: 02FEB2007

TEST	SPECIFICATIONS	RESULTS
Viable Cell Count by Hemacytometry (pre-freeze)	> 10 <sup>5</sup> cells/mL	> 10 <sup>5</sup> cells/mL
Viability (post-freeze) <sup>2</sup>	Growth	Growth
Sterility (21-day incubation) Harpo's HTYE broth <sup>3</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth	No growth

NR-2595 was produced by inoculation of ATCC<sup>®</sup> 50548™ into LYI *Entamoeba* Medium (LEM; <u>ATCC medium 2154:</u> ). The culture was incubated in an axenic and microaerophilic environment for 4 days at 35°C, until peak density was reached.

**Date:** 15 OCT 2009 **Signature:** Signature on File

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

Biodefense and Emerging Infections Research Resources Repository

800-359-7370

Fax: 703-365-2898

NR-2595\_5078409\_15OCT2009

<sup>&</sup>lt;sup>2</sup>Incubated under propagation conditions for 3 days at 35°C in an axenic and microaerophilic environment.

<sup>&</sup>lt;sup>3</sup>Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.