

Certificate of Analysis for NR-2760

Influenza A Virus, A/chicken/Germany/N/49 (H10N7)

Catalog No. NR-2760

(Derived from ATCC® VR-1334™)

Product Description: Pooled allantoic fluid from specific-pathogen free (SPF) embryonated chicken eggs¹ infected with influenza A virus, A/chicken/Germany/N/49 (H10N7).

Lot²: 7677044 Manufacturing Date: 05OCT2006

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs ¹ Hemagglutination assay	Active	Active
Sequencing of Species-Specific Region	Influenza A virus	Influenza A virus
Titer by CEID₅₀ Assay³,⁴ in Embryonated Chicken Eggs¹	Report results	1.6 X 10 ⁷ CEID ₅₀ /mL
RT-PCR Assay of Extracted RNA ⁵	~ 1030 bp amplicon	~ 1030 bp amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹10-day-old SPF Fertile Chicken Eggs acquired from B&E Eggs, York Springs, PA.

Date: 10 OCT 2007 **Signature:** Signature on File

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

© 2007 American Type Culture Collection (ATCC). All rights reserved.

Page 1 of 1

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

Biodefense and Emerging Infections Research Resources Repository P.O. Box 4137 Manassas, VA 20108-4137 USA

www.beiresources.org

Fax: 703-365-2898

800-359-7370

²NR-2760 was produced by growth of ATCC[®] VR-1334[™] (Lot 214343) in the allantoic cavity of embryonated chicken eggs¹ for 2 days at 35°C in a humidified chamber without CO₂.

 $^{^3}$ The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁴48 hours at 35°C in a humidified chamber without CO₂.

⁵BM-M1 and BM-M-1027R primers; Obenauer, J. C., et al. "Large-Scale Sequence Analysis of Avian Influenza Isolates." <u>Science</u> 311 (2006): 1576–1580. PubMed: 16439620.

⁶Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.