

## **Certificate of Analysis for NR-112**

## Listeria monocytogenes, Strain Li 2108

Catalog No. NR-112

(Derived from ATCC® 19117™)

**Product Description**: *Listeria monocytogenes* (*L. monocytogenes*) is a Gram-positive, facultative intracellular bacterium that is extremely tolerant of external stresses.

Lot<sup>1</sup>: 3656833 Manufacturing Date: 24MAR2004

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis	Crom positive red	Crom positive red
Cellular morphology Colony morphology <sup>2</sup>	Gram-positive rod Report results	Gram-positive rod Circular, entire, convex, gray-white
Biochemical characterization: Analytical profile index (API <sup>®</sup> Listeria) β-hemolysis <sup>3</sup> Catalase activity	Consistent with <i>L. monocytogenes</i> Report results Positive	Consistent with <i>L. monocytogenes</i> Positive Positive
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 520 bp)	Consistent with L. monocytogenes	Consistent with <i>L. monocytogenes</i> <sup>4</sup>
Viability (post-freeze) <sup>5</sup>	Growth	Growth

<sup>&</sup>lt;sup>1</sup>NR-112 was prepared by Brain Heart Infusion broth (BD 237500) culture of ATCC<sup>®</sup> 19117™ for 24 hours at 37°C and aerobic atmosphere.

www.beiresources.org

**Date:** 27 AUG 2008 **Signature:** Signature on File

**Title:** Technical Manager, BEI Authentication or designee

ATCC<sup>®</sup>, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC<sup>®</sup>'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

**Biodefense and Emerging Infections Research Resources Repository** P.O. Box 4137 Manassas, VA 20108-4137 USA

800-359-7370

<sup>&</sup>lt;sup>2</sup>24 hours at 37°C and aerobic atmosphere in Tryptic Soy Agar (BD 236950) with 5% defibrinated sheep blood

<sup>&</sup>lt;sup>3</sup>Hemolytic activity on blood agar is generally used to distinguish *L. monocytogenes* from other *Listeria* species.

<sup>&</sup>lt;sup>4</sup>Also consistent with other *Listeria* species

<sup>&</sup>lt;sup>5</sup>24 hours at 37°C and aerobic atmosphere