

Certificate of Analysis for NR-26742

Borrelia burgdorferi, Signature-Tagged Mutagenesis Library Clone T11TC515 (Gene IR_BB_P34-BB_P35)

Catalog No. NR-26742

Product Description: Borrelia burgdorferi (B. burgdorferi), strain B31 5A18NP1 STM library clone T11TC515 was produced by signature-tagged mutagenesis (STM) of the intergenic region between the BB_P34 and BB_P35 genes.

Lot¹: 70021443 Manufacturing Date: 31JAN2019

| TEST | SPECIFICATIONS | RESULTS |
|---|--------------------------------------|--|
| Phenotypic Analysis Cellular morphology ² Motility (wet mount) | Spirochete Report results | Spirochete Motile |
| Purity (post-freeze) ³ | No growth observed | No growth observed |
| Viability (post-freeze) Visual observation LIVE/DEAD [®] BacLight™ Bacterial Viability | Growth Green fluorescence visible | Growth ² Green fluorescence visible (Figure 1) ⁴ |

¹NR-26742 was produced by inoculation of the deposited material into Revised Barbour-Stoenner-Kelly broth supplemented with 200 μg/mL kanamycin and 40 μg/mL gentamicin and grown for 10 days at 32°C in a microaerophilic atmosphere to produce this lot.

Figure 1: LIVE/DEAD® BacLight™ Bacterial Viability



/Heather Couch/ Heather Couch

01 MAR 2019

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources

www.beiresources.org

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²7 days at 32°C in a microaerophilic atmosphere in Revised Barbour-Stoenner-Kelly broth supplemented with 200 μg/mL kanamycin and 40 μg/mL gentamicin

³Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere with 5% CO₂ on Tryptic Soy agar with 5% defibrinated sheep blood.

⁴Determined with LIVE/DEAD® *Bac*Light™ Bacterial Viability Kit, 100x magnification (Invitrogen™ L7007) after a 7-day incubation at 32°C in a microaerophilic atmosphere in Revised Barbour-Stoenner-Kelly broth supplemented with 200 μg/mL kanamycin and 40 μg/mL gentamicin. Cells with a compromised membrane that are dead or dying will stain red, while cells with an intact membrane will stain green.