

## **Certificate of Analysis for NR-26378**

Borrelia burgdorferi, Signature-Tagged Mutagenesis Library Clone T11TC063 (IR\_BB\_O28-BB\_O29)

Catalog No. NR-26378

**Product Description:** Borrelia burgdorferi (B. burgdorferi), strain B31 5A18NP1 STM library clone T11TC063 was produced by signature-tagged mutagenesis (STM) of the intergenic region between the BB\_O28 and BB\_O29 genes.

Lot<sup>1</sup>: 70021451 Manufacturing Date: 25JAN2019

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology <sup>2</sup> Motility (wet mount)	Spirochete Report results	Spirochete Motile
Purity (post-freeze) <sup>3</sup>	No growth observed	No growth observed
Viability (post-freeze) Visual observation LIVE/DEAD <sup>®</sup> BacLight™ Bacterial Viability	Growth Green fluorescence visible	Growth <sup>2</sup> Green fluorescence visible (Figure 1) <sup>4</sup>

<sup>&</sup>lt;sup>1</sup>NR-26378 was produced by inoculation of the deposited material into Revised Barbour-Stoenner-Kelly medium supplemented with 200 μg/mL kanamycin and 40 μg/mL gentamicin and grown for 9 days at 32°C in a microaerophilic atmosphere to produce this lot.

Figure 1: LIVE/DEAD® BacLight™ Bacterial Viability



/Heather Couch/

www.beiresources.org

Heather Couch 12 APR 2019

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

<sup>&</sup>lt;sup>2</sup>10 days at 32°C in a microaerophilic atmosphere in Revised Barbour-Stoenner-Kelly broth supplemented with 200 μg/mL kanamycin and 40 μg/mL gentamicin

³Purity of this lot was assessed for 10 days at 37°C in an aerobic atmosphere with 5% CO₂ on Tryptic Soy agar with 5% defibrinated sheep blood. 
⁴Determined with LIVE/DEAD® BacLight™ Bacterial Viability Kit, 1000x magnification (Invitrogen™ L7007) after a 10-day incubation at 32°C in a microaerophilic atmosphere in Revised Barbour-Stoenner-Kelly broth supplemented with 200 µg/mL kanamycin and 40 µg/mL gentamicin. Cells with a compromised membrane that are dead or dying will stain red, while cells with an intact membrane will stain green.