

Measles Virus, MVs/Florida.USA/37.23, Genotype D8

Catalog No. NR-60833

This reagent is the property of the U.S. Government.

Product Description:

Measles virus (MeV), MVs/Florida.USA/37.23, genotype D8 (D8) was collected by nasopharyngeal swab on September 14, 2023, and isolated on May 7, 2024, in Florida, USA. NR-60833 was produced by infecting *Chlorocebus aethiops* kidney epithelial cells expressing human Signaling Lymphocytic Activation Molecule (Vero-hSLAM; BEI Resources NR-55500™) with the deposited material and incubating in Dulbecco's Modified Eagle's Medium (ATCC® 30-2002™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 3 days at 37°C with 5% CO₂ to produce this lot.

Passage History:

Vh(2)/Vh(2) (Prior to deposit at BEI Resources/BEI Resources); Vh = Vero-hSLAM

Lot: 70078292

Manufacturing Date: 22SEP2025

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero-hSLAM Cells	Syncytia formation	Syncytia formation
Sequencing of Species-Specific Region (~ 450 nucleotides)	> 98% sequence identity with measles virus, MVs/Florida.USA/37.23, genotype D8 (GenBank: PP319645.1)	99.8% sequence identity with measles virus, MVs/Florida.USA/37.23, genotype D8 (GenBank: PP319645.1)
Next-Generation Sequencing (NGS) of Complete Genome Using Illumina® MiSeq™ Platform	≥ 98% sequence identity with measles virus genotype D8 strain MVs/Utah.USA/23.25 [D8], complete genome (GenBank: PX121931)	98.60% sequence identity with measles virus genotype D8 strain MVs/Utah.USA/23.25 [D8], complete genome (GenBank: PX121931) ¹
Titer by TCID₅₀ Assay in Vero-hSLAM Cells by Cytopathic Effect² (5 days at 37°C with 5% CO ₂)	Report results	1.6 × 10 ⁶ TCID ₅₀ /mL
Sterility test (Bact/ALERT 3D) iAST bottle (aerobic) at 32.5°C, 14-day incubation iNST bottle (anaerobic) at 32.5°C, 14-day incubation	No growth No growth	No growth No growth
Sterility (21-day incubation) Harpo's HTYE broth, 37°C and 26°C, aerobic ³ Trypticase Soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C, aerobic	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Sequence information for measles virus, MVs/Florida.USA/37.23 is not available in the NCBI database; nucleotide sequence obtained for NR-60833 lot 70078292 is 98.6% identical to measles virus genotype D8 strain MVs/Utah.USA/23.25 [D8], complete genome (GenBank: PX121931), another contemporary genotype D8 MeV strain.

²The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

³Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

/Sonia Bjorum Brower/

Sonia Bjorum Brower

10 DEC 2025

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

