

Bacteroides stercoris, Strain CC31F

Catalog No. HM-1036

Product Description:

Bacteroides stercoris (*B. stercoris*), strain CC31F was isolated in 2010 from colonic biopsy tissue of a human subject in Victoria, British Columbia, Canada. HM-1036 was produced by inoculation of BEI Resources seed lot 62344140 into Modified Reinforced Clostridial broth and grown for 2 days at 37°C in an anaerobic atmosphere (< 5% O₂; Remel™ Pack-Anaero™). Broth inoculum was added to Modified Reinforced Clostridial broth, which was grown for 2 days at 37°C in an anaerobic atmosphere to produce this lot. Quality control testing was completed under propagation conditions unless otherwise noted.


Note: Quality control of HMP material is only performed to demonstrate that the material distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited organism.

Lot: 70085372

Manufacturing Date: 04SEP2026

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology 2 days at 37°C in an anaerobic atmosphere in Tryptic Soy agar with 5% defibrinated sheep blood Colony morphology 2 days at 37°C in an anaerobic atmosphere in Tryptic Soy agar with 5% defibrinated sheep blood Motility (wet mount)	Report results Report results Report results	Gram-negative rod Circular, low convex, entire, smooth and white (Figure 1) Non-motile
Genotypic Analysis Sequencing of 16S ribosomal RNA (rRNA) gene (~ 1410 base pairs)	≥ 99% identical to <i>B. stercoris</i> , strain CC31F (GenBank: ATFP01000001.1)	99.5% identical to <i>B. stercoris</i> , strain CC31F (GenBank: ATFP01000001.1)
Purity (post-freeze) Anaerobic 7 days at 37°C in an anaerobic atmosphere in Tryptic Soy agar with 5% defibrinated sheep blood Aerobic 7 days at 37°C in an aerobic atmosphere with 5% CO ₂ in Tryptic Soy agar with 5% defibrinated sheep blood	Growth consistent with expected colony morphology Report results	Growth consistent with expected colony morphology No growth
Viability (post-freeze) 2 days at 37°C in an anaerobic atmosphere in Tryptic Soy agar with 5% defibrinated sheep blood	Growth	Growth

Figure 1: Colony Morphology



/Sonia Bjorum Brower/

Sonia Bjorum Brower

07 OCT 2026

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

