

Monoclonal Anti-Shiga Toxin 2 Subunit B, Clone 5A5-E7 (immunoglobulin G, mouse)

Catalog No. NR-10183

This reagent is the tangible property of the U.S. Government.

For research use only. Not for human use.

Contributor:

Alison D. O'Brien, Ph.D., Chairperson, and James F. Sinclair, Ph.D., Laboratory Supervisor, Department of Microbiology and Immunology, Uniformed Services University of the Health Sciences, Bethesda, Maryland

Product Description:

Antibody Class: IgG

Mouse monoclonal antibody prepared against the B subunit of Shiga-like toxin 2 from *Escherichia coli* (*E. coli*) was purified from mouse ascites by protein G affinity chromatography.

The term Shiga toxin (Stx) refers to two families of related toxins: Shiga toxin/Shiga-like toxin 1 and Shiga-like toxin 2.^{1,2} Shiga toxin is produced by *Shigella dysenteriae*, while Shiga-like toxin 1 and Shiga-like toxin 2 are both produced by enterohemorrhagic strains of *E. coli*. Stx are multimeric molecules that are comprised of two polypeptide subunits, A and B. The Stx B subunit is a pentamer that binds the toxin to glycolipids on host cell membranes and the entire Stx molecule can then enter the cell via endocytosis.³ Once inside the cell, the Stx A subunit undergoes proteolytic cleavage and the reduction of an internal disulfide bond to generate Stx A₁ and Stx A₂. Stx A₁ is an N-glycosidase that catalytically inactivates the 28S ribosomal RNA subunit to inhibit protein synthesis.⁴ The nucleotide sequences of the genes for the Shiga-like toxin 1 B subunit from *E. coli* (GenBank: AB035142)⁵ and the Stx B subunit from *S. dysenteriae* (GenBank: M24352)⁶ have been reported.

Material Provided:

Each vial contains approximately 50 µg of NR-10183 in PBS. The concentration, expressed as mg per mL, is shown on the Certificate of Analysis.

Packaging/Storage:

NR-10183 was packaged aseptically in vials. The product is provided frozen on dry ice and should be stored at -20°C or colder immediately upon arrival. Once thawed, the unused material may be stored at 4°C. Freeze-thaw cycles should be avoided.

Functional Activity:

NR-10183 is reactive with the recombinant B subunit of Shiga toxin 2 from *E. coli* as shown by ELISA.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: Monoclonal Anti-Shiga Toxin 2 Subunit B, Clone 5A5-E7, NR-10183."

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmbl5/bmbl5toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Sandvig, K. "Shiga Toxins." Toxicon 39 (2001): 1629–1635. PubMed: 11595626.

2. O'Loughlin, E. V. and R. M. Robins-Browne. "Effect of Shiga Toxin and Shiga-like Toxins on Eukaryotic Cells." Microbes Infect. 3 (2001): 493–507. PubMed: 11377211.
3. Sandvig, K., et al. "Endocytosis from Coated Pits of Shiga Toxin: A Glycolipid-binding Protein from *Shigella dysenteriae* 1." J. Cell Biol. 108 (1989): 1331–1343. PubMed: 2564398.
4. Skinner, L. M. and M. P. Jackson. "Investigation of Ribosome Binding by the Shiga Toxin A1 Subunit, Using Competition and Site-Directed Mutagenesis." J. Bacteriol. 179 (1997): 1368–1374. PubMed: 9023224.
5. Yokoyama, S., et al. "Nucleotide Sequence Analysis of Shiga (-like) Toxins from an Enterohemorrhagic *Escherichia coli* Isolated from Gifu Outbreak." Unpublished. GenBank: AB035142.
6. Kozlov, Yu. V., et al. "The Primary Structure of the Operons Coding for *Shigella dysenteriae* Toxin and Temperature Phage H30 Shiga-like Toxin." Gene 67 (1988): 213–221. PubMed: 3049254. GenBank: M24352.

ATCC® is a trademark of the American Type Culture Collection.

