

***Burkholderia thailandensis*, Strain DW503**

Catalog No. NR-4075

For research use only. Not for human use.

Contributor:

Herbert P. Schweizer, Ph.D., Department of Microbiology, Immunology and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, Colorado

Product Description:

Bacteria Classification: *Burkholderiaceae*, *Burkholderia*

Species: *Burkholderia thailandensis* (formerly *Burkholderia pseudomallei*-like or *Burkholderia pseudomallei*, Ara⁺ Biotype)^{1,2}

Strain: DW503

Original Source:³⁻⁶ *Burkholderia thailandensis* (*B. thailandensis*), strain DW503 is an allelic exchange strain of an environmental isolate, strain E264 (type strain for *B. thailandensis*), which was isolated from a rice field soil sample in central Thailand.

Comment: The entire genome sequence of *B. thailandensis*, strain E264 has been sequenced (GenBank: CP000085 and CP000086).⁷

B. thailandensis are saprophytic motile, aerobic, Gram-negative coccobacilli. *B. thailandensis* is genetically similar to both *B. mallei* and *B. pseudomallei* but lacks at least one pathogenicity island and is an avirulent species. In addition to its avirulence it can be differentiated from *B. pseudomallei* by some or all of the following: biochemical differences (assimilation of L-arabinose, 5-keto-gluconate, and adonitol, and no utilization of erythritol and dulcitol); differences in the 16S sequence (15 nucleotide dissimilarities); differences in lipopolysaccharide composition; and colony morphology on Ashdown's selective media.^{1,2} *B. thailandensis* is commonly found in Southeast Asia (central Thailand in particular) and some isolates have been obtained from northern Australia.⁸ Typical *B. thailandensis* are resistant to aminoglycosides but sensitive to tetracycline, ceftazidime and trimethoprim.¹

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in LB Broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-4075 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -80°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Luria Bertani (LB) Broth

LB Agar

Incubation:

Temperature: 30 or 37°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; thaw slowly.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tubes and plate at 30 or 37°C for 48 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Burkholderia thailandensis*, Strain DW503, NR-4075."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see www.cdc.gov/od/ohs/biosfty/bmbl5/bmbl5toc.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. Brett, P. J., D. Deshazer and D. E. Woods. "*Burkholderia thailandensis* Sp. Nov., a *Burkholderia pseudomallei*-Like Species." Int. J. Syst. Bacteriol. 48 Pt 1 (1998): 317-320. PubMed: 9542103.
2. Woods, D. E. "Species versus Biotype Status." J. Clin. Microbiol. 37 (1999): 3786-3787. PubMed: 10610379.
3. Burtnick, M., et al. "Identification of the Acid Phosphatase (*acpA*) Gene Homologues in Pathogenic and Non-Pathogenic *Burkholderia* Spp. Facilitates *TnphoA* Mutagenesis." Microbiology. 147 (2001): 111-120. PubMed: 11160805.
4. Brett, P. J., D. Deshazer and D. E. Woods. "Characterization of *Burkholderia pseudomallei* and *Burkholderia pseudomallei*-Like Strains." Epidemiol. Infect. 118 (1997): 137-148. PubMed: 9129590.
5. Ulrich, R. L., et al. "Mutational Analysis and Biochemical Characterization of the *Burkholderia thailandensis* DW503 Quorum-Sensing Network." J. Bacteriol. 186 (2004): 4350-4360. PubMed: 15205437.
6. Wuthiekanun, V., et al. "Biochemical Characteristics of Clinical and Environmental Isolates of *Burkholderia pseudomallei*." J. Med. Microbiol. 45 (1996): 408-412. PubMed: 8958243.
7. Kim, H. S., et al. "Bacterial Genome Adaptation to Niches: Divergence of the Potential Virulence Genes in Three *Burkholderia* Species of Different Survival Strategies." BMC Genomics. 6 (2005): 174. PubMed: 16336651. GenBank: CP000085 and CP000086.
8. Gee, J. E., et al. "Recovery of a *Burkholderia thailandensis*-Like Isolate from an Australian Water Source." BMC Microbiol. 8 (2008): 54. PubMed: 18384685.
9. Inglis, T. J., et al. "Cellular Fatty Acid Profile Distinguishes *Burkholderia pseudomallei* from Avirulent *Burkholderia thailandensis*." J. Clin. Microbiol. 41 (2003): 4812-4814. PubMed: 14532228.
10. Inglis, T. J., et al. "Comparison of Diagnostic Laboratory Methods for Identification of *Burkholderia pseudomallei*." J. Clin. Microbiol. 43 (2005): 2201-2206. PubMed: 15872242.

ATCC® is a trademark of the American Type Culture Collection.

