

## *Giardia lamblia*, Strain Sug

Catalog No. NR-9233

**For research use only. Not for human use.**

### Contributor:

Dr. Theodore E. Nash, Chief, Gastrointestinal Parasites Section, Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland

### Product Description:

Protozoa Classification: Hexamitidae, Giardiinae, *Giardia*

Species: *Giardia lamblia* (also referred to as *Giardia intestinalis* and *Giardia duodenalis*)

Strain: Sug

Original Source: *Giardia lamblia* (*G. lamblia*) strain Sug was isolated from a human in Maryland, U. S. A.<sup>1</sup>

Comments: Genome sequencing information for *G. lamblia* is available at [Giardia lamblia Genome Project](#).<sup>2,3</sup>

*G. lamblia* is a pear-shaped, flagellated protozoan that causes a wide variety of gastrointestinal complaints and is one of the most common causes of parasite infection of humans worldwide, and the second most common in the United States. The disease is commonly water-borne because *Giardia* cysts are resistant to the chlorine levels in normal tap water and survive well in cold mountain streams. Food-borne transmission is rare but can occur with ingestion of raw or undercooked foods. Giardiasis is a zoonosis, and cross-infectivity among beaver, cattle, dogs, rodents, and bighorn sheep provides a constant reservoir. The life cycle of *Giardia* consists of two stages: the fecal-orally transmitted cyst and the disease-causing trophozoite. Cysts are passed in a host's feces, remaining viable in a moist environment for months. Ingestion of 10 to 25 cysts can cause infection in humans.<sup>4</sup>

### Material Provided:

Each vial of NR-9233 contains approximately 0.5 mL of culture in cryopreservative. Please see Appendix I below for cryopreservation instructions.

### Packaging/Storage:

NR-9233 was packaged aseptically in screw-capped plastic cryovials and is provided frozen on dry ice. The product should be stored at cryogenic temperature (-130°C or colder), preferably in the vapor phase of a liquid nitrogen freezer. If liquid nitrogen storage facilities are not available, frozen cryovials may be stored at -70°C or colder for approximately one week. Note: Do not under any circumstances store vials at temperatures warmer than -70°C. Storage under these conditions will result in the death of the culture.

To insure the highest level of viability, the culture should be

initiated immediately upon receipt. Any warming of the product during shipping and transfer must be avoided, as this will adversely affect the viability of the product. For transfer between freezers and for shipping, the product may be placed on dry ice for brief periods, although use of a portable liquid nitrogen carrier is preferred. Please read the following recommendations prior to using this material.

### Growth Conditions:

#### Growth Media:

ATCC® medium 2695 (previously ATCC® medium 1404), supplemented with Diamond's Vitamin Solution and 10% heat-inactivated adult bovine serum. Please see Appendix II for media preparation instructions.

Note: An alternative to ATCC® medium 2695 is ATCC® medium PRA-2155.

#### Incubation:

Temperature: 35°C

Atmosphere: axenic and microaerophilic

#### Propagation:

1. Thaw frozen ampoule in a 35°C water bath, for 2 to 3 min. Immerse the ampoule just sufficient to cover the frozen material. Do not agitate the ampoule.
2. Immediately after thawing, aseptically transfer contents to a 16 x 125 mm screw-capped borosilicate glass test tube containing 13 mL of ATCC® medium 2695. Incubate the tube on a 15° horizontal slant at 35°C.

#### Maintenance:

1. When the culture has reached or is near peak density, place the tubes on ice for 10 minutes.
2. Gently invert the culture tube 10 times and aseptically transfer a 0.1 to 0.4 mL aliquot to a screw-capped test tube containing 13 mL ATCC® medium 2695.
3. Incubate the culture on a 15° horizontal slant at 35°C.
4. Transfer the culture every 3 to 4 days as described in Maintenance steps 1 and 2. The transfer interval will depend on the size of the inoculum and the quality of the medium. This should be determined empirically by examining the culture on a daily basis until conditions for stable growth have been achieved. Do not allow the culture to overgrow. Viability of the culture may be affected soon after reaching peak density.

### Citation:

Acknowledgment for publications should read "The following reagent was obtained through the NIH Biodefense and Emerging Infections Research Resources Repository, NIAID, NIH: *Giardia lamblia*, Strain Sug, NR-9233."

### Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2007; see

[www.cdc.gov/od/ohs/biosfty/bmb15/bmb15toc.htm](http://www.cdc.gov/od/ohs/biosfty/bmb15/bmb15toc.htm).

### Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at [www.beiresources.org](http://www.beiresources.org).

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government make any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

### Use Restrictions:

**This material is distributed for internal research, non-commercial purposes only.** This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

### References:

1. Nash, T. E. and D. B. Keister. "Differences in Excretory-Secretory Products and Surface Antigens among 19 Isolates of *Giardia*." J. Infect. Dis. 152 (1985): 1166-1171. PubMed: 4067331.
2. Morrison, H. G., et al. "Genomic Minimalism in the Early Diverging Intestinal Parasite *Giardia lamblia*." Science 317 (2007): 1921-1926. PubMed: 17901334.
3. McArthur, A. G., et al. "The *Giardia* Genome Project Database." FEMS Microbiol. Lett. 189 (2000): 271-273. PubMed: 10930750.
4. Kucik, C. J., G. L. Martin, and B. V. Sorter. "Common Intestinal Parasites." Am. Fam. Physician 69 (2004): 1161-1168. PubMed: 15023017.

ATCC® is a trademark of the American Type Culture Collection.



### APPENDIX I: CRYOPRESERVATION

1. Harvest cells from a culture that is at or near peak density. To detach cells from the wall of the culture tubes place on ice for 10 min. Invert tubes several times until the majority of the cells are in suspension. Centrifuge tubes at 800 x g for 5 minutes.
2. Adjust the concentration of cells to  $1-2 \times 10^7$ /mL in fresh medium.
3. Before centrifuging prepare a 24% (v/v) solution of sterile DMSO in fresh medium containing 8% (w/v) sucrose. The solution is prepared as follows:
  - a) Add 1.05 g sucrose to 10 mL of fresh medium and filter sterilize through a 0.2  $\mu$ m filter
  - b) Add 2.4 mL of DMSO to an ice cold 20 x 150 mm screw-capped test tube
  - c) Place the tube on ice and allow the DMSO to solidify (~ 5 min) and then add 7.6 mL of ice cold medium prepared in step 3a. The final concentration will be 24% (v/v) DMSO and 8% (w/v) sucrose
  - d) Invert several times to dissolve the DMSO
  - e) Allow to warm to room temperature.
4. Mix the cell preparation and the cryoprotective agent, prepared in step 3, in equal portions. Thus, the final concentration will equal 12% (v/v) DMSO, 4% (w/v) sucrose and  $10^7$  cells/mL. **The time from the mixing of the cell preparation and DMSO stock solution before the freezing process is begun should be no less than 15 min. and no longer than 30 min.**
5. Dispense in 0.5 mL aliquots into 1.0 - 2.0 mL sterile plastic screw-capped cryovials.
6. Place the vials in a controlled rate freezing unit. From room temperature, cool at  $-1^\circ\text{C}/\text{min}$  to  $-40^\circ\text{C}$ . If the freezing unit can compensate for the heat of fusion, maintain rate at  $-1^\circ\text{C}/\text{min}$  through the heat of fusion. At  $-40^\circ\text{C}$ , plunge into liquid nitrogen. Alternatively, place the vials in a Nalgene  $1^\circ\text{C}$  freezing apparatus. Place the apparatus at  $-80^\circ\text{C}$  for 1.5 to 2 hours and then plunge ampoules into liquid nitrogen (The cooling rate in this apparatus is approximately  $-1^\circ\text{C}/\text{min}$ ).
7. The frozen preparations should be stored in either the vapor or liquid phase of a nitrogen freezer. Stability of frozen preparations is higher at  $-130^\circ\text{C}$  as compared to temperatures warmer than  $-130^\circ\text{C}$ .

**APPENDIX II: MEDIA**

**ATCC® Medium 2695 (Keister's Modified TYI-S-33)**

Casein Digest (BD Trypticase 211705)	20.0 g
Yeast Extract (BD 212750)	10.0 g
Bovine Bile (Sigma B-8381)	0.75 g
NaCl	2.0 g
L-Cysteine HCl	2.0 g
Ascorbic Acid	0.2 g
K <sub>2</sub> HPO <sub>4</sub>	1.0 g
KH <sub>2</sub> PO <sub>4</sub>	0.6 g
Ferric Ammonium Citrate	22.8 mg
Distilled water	880 mL

1. Dissolve the ingredients in 880 mL of distilled water in the order indicated.
2. Adjust the pH to 7.0 to 7.2 with 1 N NaOH.
3. Filter sterilize.
4. To prepare the complete medium, aseptically add 20 mL of Diamond's Vitamins Solution and 100 mL of heat-inactivated adult bovine serum.
5. Mix thoroughly and distribute 13 mL aliquots into 16 x 125 mm screw-capped borosilicate glass test tubes. Store at 4 to 8°C in the dark. Use within 7 to 10 days. Long term storage may result in the formation of precipitates and failure to support growth of *Giardia*.

**NOTE:** Serum is heat-inactivated by exposure to 56°C for 30 minutes to inactivate proteins of the complement pathway.

**Diamond's Vitamin Solution**

**Solution 1:** (DL-6,8-Thioctic acid [DL-α-Lipoic acid], 1mg/mL).

Dissolve 100 mg of DL-6,8-Thioctic acid (oxidized form, Sigma T1395) in 100 mL of absolute ethanol.

**Solution 2:** (Vitamin B<sub>12</sub>, 0.4 mg/mL).

Dissolve 40 mg of vitamin B<sub>12</sub> (Sigma V2876) in 100 mL distilled water.

**Solution 3:** (Tween 80, 50% w/v).

Dissolve 50 g of Tween 80 (Sigma P1754) in 100 mL absolute ethanol.

**Solution 4 (All components from Sigma):**

α-tocopherol phosphate, disodium salt	0.025 mg
d-biotin	0.025 mg
Calciferol (Vitamin D <sub>2</sub> )	0.250 mg
Calcium D-(+)pantothenate	0.025 mg
Choline chloride	1.250 mg
Folic acid	0.025 mg
i-Inositol	0.125 mg
Menadione (Vitamin K <sub>3</sub> )	0.025 mg
Niacin	0.0625 mg
Niacinamide	0.0625 mg
p-aminobenzoic acid	0.125 mg
Pyridoxal HCl	0.0625 mg
Pyridoxine HCl	0.0625 mg
Riboflavin	0.0625 mg
Thiamine HCl	0.025 mg
Vitamin A	0.250 mg
Distilled water	1 L

To prepare Diamond's Vitamin Solution combine the following:

Solution 1	0.4 mL
Solution 2	1.2 mL
Solution 3	0.4 mL
Solution 4	100.0 mL
Sterile distilled water	18.0 mL

Filter sterilize. Store at -20°C.