

***Yersinia pestis*, Strain K25 Derivative 80 (D80)**

**Catalog No. NR-4703**

**Product Description:** *Yersinia pestis* (*Y. pestis*) is an aerobic, non-spore-forming, Gram-negative rod-shaped bacterium. *Y. pestis*, strain K25(D80) is an avirulent derivative of the K25 strain that contains the pMT1 and pPCP1 plasmids, but lacks the pCD1 plasmid that is essential for virulence as well as the unstable chromosomal *pgm* locus.

**Lot<sup>1</sup>: 58324518**

**Manufacturing Date: 19SEP2008**

TEST	SPECIFICATIONS	RESULTS
<b>Phenotypic Analysis</b> Cellular morphology Colony morphology <sup>2</sup>  Congo red (CR) agar <sup>3,4</sup> Biochemical Analyses Analytical profile index (API 20 E <sup>®</sup> ) Nitrate reduction Fermentation of glycerol Urease	Gram-negative rods Report results  No red colonies (crb <sup>-</sup> )  Consistent with <i>Y. pestis</i> Negative Positive Negative	Gram-negative rods Circular, convex, entire, opaque (Figure 1) No red colonies (crb <sup>-</sup> )  Consistent with <i>Y. pestis</i> Negative Positive Negative
<b>Genotypic Analysis</b> Sequencing of 16S ribosomal RNA gene (~ 1440 bp)	Consistent with <i>Y. pestis</i> <sup>5</sup>	Consistent with <i>Y. pestis</i> <sup>5</sup>
<b>PCR Assay of Extracted DNA</b> <i>Y. pestis</i> specific sequence (YPO0396) <sup>6</sup> 16S ribosomal RNA gene Presence of virulence-associated plasmids pMT1 (pFra; 100 kb plasmid) pCD1 (pYV; 70 kb plasmid) pPCP1 (pPla; 9.5 kb plasmid)	~ 800 bp amplicon ~ 1500 bp amplicon  ~ 1200 bp amplicon None detected ~ 400 bp amplicon	~ 800 bp amplicon ~ 1500 bp amplicon  ~ 1200 bp amplicon None detected ~ 400 bp amplicon
<b>Viability (post-freeze)<sup>2</sup></b>	Growth on agar	Growth on agar

<sup>1</sup>*Y. pestis*, strain K25(D80) was deposited by Professor Robert R. Brubaker of the Department of Microbiology and Molecular Genetics at Michigan State University, East Lansing, Michigan. NR-4703 was prepared by Tryptic Soy Broth culture of the deposited material and grown 48 hours at 28°C and aerobic atmosphere. Broth inoculum was added to Kolles which were grown 48 hours at 28°C and aerobic atmosphere to produce this lot.

<sup>2</sup>48 hours at 28°C and aerobic atmosphere on Tryptic Soy Agar

<sup>3</sup>7 days at 28°C and aerobic atmosphere on CR agar

<sup>4</sup>Hare, J. M. and K. A. McDonough. "High-Frequency RecA-Dependent and -Independent Mechanisms of Congo Red Binding Mutations in *Yersinia pestis*." *J. Bacteriol.* 181 (1999): 4896-4904. PubMed: 10438760.

<sup>5</sup>Also consistent with other *Yersinia* species

<sup>6</sup>Sequence locus tag YPO0396 codes for an uncharacterized protein that is highly conserved in *Y. pestis*

**Figure 1**



**Date:** 09 MAR 2011

**Signature:**



**Title:** Technical Manager, BEI Authentication or designee

ATCC<sup>®</sup>, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC<sup>®</sup>'s knowledge.

ATCC<sup>®</sup> is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

