

Dengue Virus Type 2, 328298

Catalog No. NR-12217

Product Description: Cell lysate and supernatant from *Aedes albopictus* clone C6/36 cells¹ infected with dengue virus type 2 (DEN-2), 328298.

Lot²: 58526834

Manufacturing Date: 27MAY2009

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in C6/36 Cells ¹	Report results	Cell rounding
Identification by Indirect Fluorescent Antibody (IFA) Assay ³	Fluorescence observed	Fluorescence observed
Sequencing of DEN-2 Specific Sequence (793 nucleotides)	Consistent with DEN-2	Consistent with DEN-2
Titer by TCID ₅₀ Assay in C6/36 Cells With IFA Readout ^{1,4,5}	Report results	1.6 x 10 ⁷ TCID ₅₀ /mL
Functional Activity by RT-PCR Assay Using DEN-2 Specific Primers	~ 1200 bp amplicon	~ 1200 bp amplicon
Bacterial Sterility (Bact/ALERT[®] 3D Microbial Detection System) 14-day incubation of NR-12217: i NST culture bottle, 32°C, anaerobic i AST culture bottle, 32°C, aerobic	No growth No growth	No growth No growth
Fungal Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C and 26°C, aerobic	No growth No growth No growth	No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

¹*Aedes albopictus* clone C6/36 cells (ATCC[®] CRL-1660[™])

²DEN-2, 328298 was deposited by Dr. Rebeca Rico-Hesse of the Department of Virology and Immunology, Southwest Foundation for Biomedical Research, San Antonio, Texas. NR-12217 was grown from deposited virus seed in Minimum Essential Medium containing Earle's salts and non-essential amino acids (Invitrogen[™] 10370-021) supplemented with 2% irradiated fetal bovine serum (Cambrex[®] 14-471F), 2 mM L-glutamine (Invitrogen[™] 25030-081), and 1 mM sodium pyruvate (Invitrogen[™] 11360-070) for 8 days at 28°C with 5% CO₂

³Using monoclonal antibody specific to DEN-2 (Chemicon MAB8702)

⁴The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁵7 days at 28°C with 5% CO₂

⁶Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 28 FEB 2011

Signature:



Title:

Technical Manager, BEI Authentication or designee

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

