

Certificate of Analysis for HM-133

Gardnerella vaginalis, Strain UPII 315-A

Catalog No. HM-133

Product Description: Gardnerella vaginalis (G. vaginalis), strain UPII 315-A was isolated from

human vaginal flora.

Lot¹: 59576255 Manufacturing Date: 09DEC2010

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ³	Gram-variable rods ² Report results	Gram-variable rods Circular and glistening (Figure 1)
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1420 base pairs)	≥ 99% identical to depositor's sequence Consistent with <i>G. vaginalis</i>	Pending Consistent with <i>G. vaginalis</i>
Viability (post-freeze) ³	Growth	Growth

G. vaginalis, strain UPII 315-A was deposited by Sharon L. Hillier, Professor, Department of Obstetrics, Gynecology and Reproductive Sciences, Magee-Womens Research Institute, University of Pittsburgh, Pittsburgh, Pennsylvania. The deposited material was inoculated into NYC III broth (ATCC medium 1685) and incubated for 24 hours at 37°C in an aerobic atmosphere with 5% CO₂. Broth inoculum was added to Kolles which were grown 48 hours at 37°C to produce this lot.

Figure 1



Date: 14 APR 2011

Signature:

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by ATCC® to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

BEI Resources

²G. vaginalis is often described as a Gram-variable organism but has a thin, Gram-positive cell wall [see Harper, J. J. and G. H. G. Davis. "Cell Wall Analysis of Gardnerella vaginalis (Haemophilus vaginalis)." Int. J. Syst. Bacteriol. 32 (1982): 48-50.]
³48 hours at 37°C and 5% CO₂ on Chocolate Agar (GC Medium)