

Bovine Parainfluenza Virus 3, SF-4

Catalog No. NR-3234

This reagent is the property of the U.S. Government.

For research use only. Not for human use.

Contributor:

National Institutes of Allergy and Infectious Disease, National Institutes of Health

Manufacturer:

Lederle Laboratories, NIH-69-2043

Product Description:

Reagent: Seed Virus

Virus Classification: *Paramyxoviridae, Respirovirus*

Agent: Bovine parainfluenza virus 3

Strain/Isolate: SF-4

NIAID Class: Research Reference Reagent

Donor: ATCC® VR-281

Donor Passage History (# of passages):

Bovine embryo kidney (12)

Producer Passage History (# of passages):

African green monkey kidney (3)

Note: BEI Resources was asked to distribute this virus preparation from NIAID's historical repository. Historical characterization information is shown below in the Functional Activity and Purity sections (tests performed in August 1970). Recent characterization information is shown on the Certificate of Analysis.

Material Provided/Storage:

Composition: Basal Medium Eagle with 4% sorbitol and 4% N-Z amine

Volume: 1.0 mL

Storage Temperature: -60°C or colder

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Functional Activity (August 1970):

Infectivity:

Conditions: African green monkey kidney

TCID₅₀:¹ 1.6 x 10⁷ per mL

Purity (August 1970):

Serum Neutralization Breakthrough: Negative

Bacterial Sterility: Negative

Mycoplasma: Negative

Citation:

Acknowledgment for publications should read "The following

reagent was obtained through BEI Resources, NIAID, NIH: Bovine Parainfluenza Virus 3, SF-4, NR-3234."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmb15/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in tissue culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the cultures inoculated, just as a Lethal Dose 50% (LD₅₀)

is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

ATCC® is a trademark of the American Type Culture Collection.

