

Certificate of Analysis for NR-540

La Crosse Virus

Catalog No. NR-540

(Derived from ATCC® VR-744™)

Product Description: Cell lysate and supernatant from African green monkey kidney (Vero) cells¹ infected with La Crosse virus.

Lot²: 3845462 Manufacturing Date: 03DEC2004

TEST	SPECIFICATIONS	RESULTS
Titer by TCID ₅₀ Assay ^{3,4} in Vero Cells ¹	Report results	6.3 X 10 ⁷ TCID ₅₀ /mL
Sterility (21-day incubation)		
Harpo's HTYE broth⁵, 37°Ć and 26°C, aerobic	No growth	No growth
Trypticase soy broth, 37°C and 26°C, aerobic	No growth	No growth
Sabouraud broth, 37°C and 26°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, anaerobic	No growth	No growth
Thioglycollate broth, 37°C, anaerobic	No growth	No growth
DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination		
Agar and broth culture (14-day incubation at 37°C)	None detected	None detected
DNA Detection by PCR of extracted Test Article nucleic acid	None detected	None detected

¹Vero cells: ATCC[®] CCL-81™.

Date: 18 OCT 2006 **Signature:** Signature on File

Title: Technical Manager, BEI Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

Fax: 703-365-2898

E-mail: contact@beiresources.org

800-359-7370

²Grown in Eagle's Minimum Essential Medium containing Earle's salts and non-essential amino acids (ATCC[®] 30-2003) supplemented with 2% fetal bovine serum (ATCC[®] 30-2020) for 2 days at 37°C with 5% CO₂.

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴7 days at 37°C and 5% CO₂.

⁵Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.