

Certificate of Analysis for NR-41802

Influenza B Virus, B/Hong Kong/330/2001 (Victoria Lineage)

Catalog No. NR-41802

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs¹ infected with influenza B virus, B/Hong Kong/330/2001 (Victoria Lineage)

Lot^{2,3}: 61361421 Manufacturing Date: 29NOV2012

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs ¹ Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin Coding Region (982 nucleotides)	Consistent with B/Hong Kong/330/2001 (Victoria Lineage)	100% identity with B/Hong Kong/330/2001 (Victoria Lineage) (Gen Bank: CY018709)
Titer by CEID ₅₀ Assay ^{4,5} in Embryonated Chicken Eggs ¹	Report results	1.6 × 10 ⁸ CEID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹⁰⁻day-old SPF Embryonated Chicken Eggs acquired from B&E Eggs, York Springs, Pennsylvania

Date: 06 MAY 2013

Signature:

Title:

Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC® s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org Tel: 800-359-7370

Fax: 703-365-2898

²Derived from CDC ID No. 2001701830

³Grown in the allantoic cavity of embryonated chicken eggs¹ for 2 days at 34°C in a humidified chamber

⁴The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁵2 days at 34°C in a humidified chamber

⁶Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.