

Certificate of Analysis for HM-665

Bacteriophage Phi_TK1932sp/Pa1651, Infectious for Pseudomonas aeruginosa

Catalog No. HM-665

Product Description: Bacteriophage phi_TK1932sp/Pa1651 was isolated from TK1932 bacterium cultured from leg, foot or decubitus ulcers of a human patient. Bacteriophage phi_TK1932sp/Pa1651 (HMP ID 9775) is a reference genome for The Human Microbiome Project (HMP).

Lot^{1,2}: 59978218 Manufacturing Date: 02JUN2011

TEST	SPECIFICATIONS	RESULTS
Plaque Morphology ³	Report results	Circular
Titer ³ Plaque-forming units (pfu) with Pseudomonas aeruginosa	>10 ⁶ pfu per mL	9 x 10 ⁹ pfu per mL
Sterility	0.22 µm filtered	0.22 µm filtered
Bacterial Inactivation 10% of total yield plated on Tryptic Soy Agar with 5% defibrinated sheep blood ⁴	No viable bacteria detected	No viable bacteria detected

Quality control of HMP material is only performed to demonstrate that the product distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited material.

⁴7 days at 37°C in an aerobic atmosphere

Date: 09 APR 2013

Signature:

Title: Technical Manager, BEI Authentication or designee

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

BEI Resources

²The deposited material was added to a Luria-Burtani (LB) soft agar overlay (0.5%) supplemented with 10 mM MgSO₄ containing the phage host, *Pseudomonas aeruginosa*, strain Pa1651. The soft agar overlay was added to an LB agar Kolle supplemented with 10 mM MgSO₄ and incubated 24 hours at 30°C in an aerobic atmosphere. After two additional passages under the above conditions, the bacteriophage were harvested to produce this lot.

³Lysis was observed on the soft agar overlay after 24 hours at 30°C in an aerobic atmosphere.