

Certificate of Analysis for HM-658

Bacteriophage Phi_MX560sp/Pa1651, Infectious for Pseudomonas aeruginosa

Catalog No. HM-658

Product Description: Bacteriophage phi_MX560sp/Pa1651 was isolated from MX560 bacterium cultured from leg, foot or decubitus ulcers of a human patient. Bacteriophage phi_MX560sp/Pa1651 (HMP ID 9768) is a reference genome for The Human Microbiome Project (HMP).

Lot^{1,2}: 59978210 Manufacturing Date: 29JUN2011

TEST	SPECIFICATIONS	RESULTS
Plaque Morphology ³	Report results	Circular and clear
Titer³ Plaque-forming units (pfu) with <i>Pseudomonas aeruginosa</i>	>10 ⁶ pfu per mL	2 x 10 ¹⁰ pfu per mL
Sterility	0.22 µm filtered	0.22 µm filtered
Bacterial Inactivation 10% of total yield plated on Tryptic Soy Agar with 5% defibrinated sheep blood ⁴	No viable bacteria detected	No viable bacteria detected

¹Quality control of HMP material is only performed to demonstrate that the product distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited material.

⁴7 days at 37°C in an aerobic atmosphere

Date: 09 APR 2013

Signature:

Title: Technical Manager, BEI Authentication or designee

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

E-mail: contact@beiresources.org

Tel: 800-359-7370 Fax: 703-365-2898

²The deposited material was added to a Luria-Burtani (LB) soft agar overlay (0.5%) supplemented with 10 mM MgSO₄ containing the phage host, *Pseudomonas aeruginosa*, strain Pa1651. The soft agar overlay was added to LB Agar Kolles supplemented with 10 mM MgSO₄ and incubated 24 hours at 30°C in an aerobic atmosphere. After three additional passages under the above conditions, the bacteriophage were harvested to produce this lot.

³Lysis was observed on the soft agar overlay after 24 hours at 30°C in an aerobic atmosphere.