

### ***Vibrio parahaemolyticus*, Isolate T6**

**Catalog No. NR-34578**

**For research use only. Not for human use.**

**Contributor:**

T. Madigan (Senior Research Officer) and D. May (Research Scientist), The Southern Australian Research and Development Institute (SARDI), Adelaide, South Australia

**Manufacturer:**

BEI Resources

**Product Description:**

Bacteria Classification: *Vibrionaceae*, *Vibrio*

Species: *Vibrio parahaemolyticus*

Isolate: T6

Original Source: *Vibrio parahaemolyticus* (*V. parahaemolyticus*), isolate T6 was obtained from an oyster in New South Wales, Australia, 2010.<sup>1</sup>

Comments: Isolate T6 was identified by growth on TCBS (Thiosulfate-Citrate-Bile-Sucrose) agar and PCR analyses.<sup>1</sup>

*V. parahaemolyticus* is a halophilic, Gram-negative motile, curved-rod shaped bacterium with a single polar flagellum. It is found in estuarine and coastal waters worldwide (Spain, Asia, Russia, South America, Africa and the United States). It is the leading cause of foodborne gastroenteritis. It is usually ingested in undercooked or raw seafood.<sup>2</sup>

*V. parahaemolyticus* is serotyped on the basis of somatic (O) and capsular (K) antigens, and is classified into at least 11 O-serogroups<sup>3</sup> and over 70 K-serogroups. Certain serotypes, including O3:K6, O1:KUT, O4:K12 and O4:K68, have been reported to be more virulent<sup>2,3</sup> and are considered to be the dominant serotypes responsible for infection.

**Material Provided:**

Each vial contains approximately 0.5 mL of bacterial culture in Tryptic Soy broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

**Packaging/Storage:**

NR-34578 was packaged aseptically in cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

**Growth Conditions:**

Media:

Tryptic Soy broth or equivalent

Tryptic Soy agar with 5% defibrinated sheep blood or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use; then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate at 37°C for 24 hours.

**Citation:**

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Vibrio parahaemolyticus*, Isolate T6, NR-34578."

**Biosafety Level: 2**

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see [www.cdc.gov/biosafety/publications/bmbl5/index.htm](http://www.cdc.gov/biosafety/publications/bmbl5/index.htm).

**Disclaimers:**

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at [www.beiresources.org](http://www.beiresources.org).

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither the ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

# Use Restrictions:

**This material is distributed for internal research, non-commercial purposes only.** This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

# References:

1. T. Madigan and D. May, Personal Communication.
2. Jones, J. L., et al. "Biochemical, Serological, and Virulence Characterization of Clinical and Oyster *Vibrio parahaemolyticus* Isolates." *J. Clin. Microbiol.* 50 (2012): 2343-2352. PubMed: 22535979.
3. Chen, M. et al. "Development of O-Serogroup Specific PCR Assay for Detection and Identification of *Vibrio parahaemolyticus*." *Int. J. Food Microbiol.* 159 (2012): 122-129. PubMed: 23072697.

ATCC® is a trademark of the American Type Culture Collection.

