

Certificate of Analysis for NR-43286

Porcine Respiratory Coronavirus, ISU-1

Catalog No. NR-43286

Derived from BEI Resources NR-448

Product Description: Clarified cell lysate and supernatant from *sus scrofa* testicular fibroblasts¹ infected with porcine respiratory coronavirus (PRCV), ISU-1

Lot²: 61617361 Manufacturing Date: 10JUL2013

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in ST Cells ¹	Cell rounding and detachment	Cell rounding and detachment
Sequencing of Species-Specific Region (410 nucleotides)	Consistent with PRCV, ISU-1	99% identity with PRCV, ISU-1 (GenBank: DQ811787)
Titer by TCID₅₀ Assay³,⁴ in ST Cells¹	Report results	8.9×10^6 TCID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

¹ST cells (ATCC[®] CRL-1746™)

Date: 21 OCT 2013

Signature:

Title:

Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²Grown in Eagle's Minimum Essential Medium (ATCC® 30-2003™) for 2 days at 37°C and 5% CO₂

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴7 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.