

***Escherichia coli*, Strain 5.3169**

Catalog No. NR-17661

For research use only. Not for human use.

Contributor:

C. DebRoy, Director, *E. coli* Reference Center, Department of Veterinary and Biomedical Sciences, College of Agricultural Sciences, The Pennsylvania State University, University Park, Pennsylvania, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Enterobacteriaceae*, *Escherichia*

Species: *Escherichia coli*

Strain: 5.3169

Serotype: O25:H4^{1,2}

Original Source: *Escherichia coli* (*E. coli*), strain 5.3169 is a human isolate from the United States, 2005.^{1,2}

Comments: *E. coli*, strain 5.3169 has been typed as an extraintestinal pathogenic *E. coli* (ExPEC) strain.^{1,2} It was selected to undergo complete genome sequencing at the [J. Craig Venter Institute](http://www.jcvi.org).

E. coli is a Gram-negative, rod-shaped bacterium commonly found in the gut flora of warm-blooded animals and is the primary facultative anaerobe of the human gastrointestinal tract. While most *E. coli* strains are harmless and are an important part of a healthy intestinal tract, some serotypes are pathogenic, causing diarrhea, urinary tract infections, respiratory illness, pneumonia, or other illnesses in their host.³⁻⁵ Pathogenic *E. coli* may be transmitted through contaminated food or water, or through contact with infected persons or animals. The six pathotypes associated with diarrhea and collectively referred to as diarrheagenic *E. coli* are: Shiga toxin-producing *E. coli* [STEC; also referred to as Verocytotoxin-producing *E. coli* (VTEC) or enterohemorrhagic *E. coli* (EHEC)]⁶, enterotoxigenic *E. coli* (ETEC)⁷, enteropathogenic *E. coli* (EPEC)⁸, enteroaggregative *E. coli* (EAEC)⁹, enteroinvasive *E. coli* (EIEC) and diffusely adherent *E. coli* (DAEC).¹⁰

ExPEC possess virulence traits that allow them to invade, colonize, and induce disease in bodily sites outside of the gastrointestinal tract. Virulence factors of ExPEC belonging to a multidrug-resistant clonal group A (CGA) have been associated with urinary tract infections.¹¹⁻¹⁴

The absence of chromosomal virulence markers *stx1* and *stx2* in NR-17661 has been confirmed by PCR amplification of extracted DNA.

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Tryptic Soy broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-17661 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy broth, Nutrient broth or equivalent
Tryptic Soy agar, Nutrient agar or Tryptic Soy agar with 5% defibrinated sheep blood or equivalent

Incubation:

Temperature: 37°C
Atmosphere: Aerobic

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read “The following reagent was obtained through BEI Resources, NIAID, NIH: *Escherichia coli*, Strain 5.3169, NR-17661.”

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmbl5/index.htm.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. DebRoy, C., Personal Communication.
2. Kapur, V., et al. "Genome Sequencing and Analysis of Pathogenic *Escherichia coli* Strains." J. Craig Venter Institute. (2009) <http://gcid.jcvi.org/docs/STEC_White_Paper.pdf>
3. Nataro, J. P. and J. B. Kaper. "Diarrheagenic *Escherichia coli*." Clin. Microbiol. Rev. 11 (1998): 142-201. PubMed: 9457432.
4. Kaper, J. B., J. P. Nataro and H. L. Mobley. "Pathogenic *Escherichia coli*." Nat. Rev. Microbiol. 2 (2004): 123-140. PubMed: 15040260.
5. Croxen, M. A., et al. "Recent Advances in Understanding Enteric Pathogenic *Escherichia coli*." Clin. Microbiol. Rev. 26 (2013): 822-880. PubMed: 24092857.
6. Smith, J. L., P. M. Fratamico and N. W. Gunther, 4th. "Shiga Toxin-Producing *Escherichia coli*." Adv. Appl. Microbiol. 86 (2014): 145-197. PubMed: 24377855.
7. Zhang, W. and D. A. Sack. "Progress and Hurdles in the Development of Vaccines against Enterotoxigenic *Escherichia coli* in Humans." Expert Rev. Vaccines 11 (2012): 677-694. PubMed: 22873126.
8. Ochoa, T. J. and C. A. Contreras. "Enteropathogenic *Escherichia coli* Infection in Children." Curr. Opin. Infect. Dis. 24 (2011): 478-483. PubMed: 21857511.
9. Estrada-Garcia, T. and F. Navarro-Garcia. "Enterotoxigenic *Escherichia coli* Pathotype: A Genetically Heterogeneous Emerging Foodborne Enteropathogen." FEMS Immunol. Med. Microbiol. 66 (2012): 281-298. PubMed: 22775224.
10. Smith, E. J., et al. "Pathogenesis of Adherent-Invasive *Escherichia coli*." Future Microbiol. 8 (2013): 1289-1300. PubMed: 24059919.
11. Johnson, J. R., et al. "Rapid and Specific Detection of *Escherichia coli* Clonal Group A by Gene-Specific PCR." J. Clin. Microbiol. 42 (2004): 2618-2622. PubMed: 15184442.
12. Johnson, J. R., et al. "A Disseminated Multidrug-resistant Clonal Group of Uropathogenic *Escherichia*

- coli* in Pyelonephritis." Lancet 359 (2002): 2249-2251. PubMed: 12103291.
13. Johnson, J. R., et al. "Distribution and Characteristics of *Escherichia coli* Clonal Group A." Emerg. Infect. Dis. 11 (2005): 141-145. PubMed: 15705341.
14. Manges, A. R., et al. "Widespread Distribution of Urinary Tract Infections Caused by a Multidrug-Resistant *Escherichia coli* Clonal Group." N. Engl. J. Med. 345 (2001): 1007-1013. PubMed: 11586952.

ATCC® is a trademark of the American Type Culture Collection.

