

**Mobala Virus, Acar 3080**

**Catalog No. NR-10173**

This reagent is the tangible property of the U.S. Government.

**Product Description:** Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells (Vero E6)<sup>1</sup> infected with Mobala virus, Acar 3080

**Lot<sup>2</sup>: 60505360**

**Manufacturing Date: 26APR2013**

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero E6 Cells <sup>1</sup> Using RT-PCR <sup>3</sup>	Report results	Infectious, no cytopathic effect
Sequencing of Species-Specific Region (230 nucleotides)	Consistent with Mobala virus, Acar 3080	100% identity with Mobala virus, Acar 3080 (GenBank: AY342390)
Titer by TCID <sub>50</sub> Assay <sup>4</sup> in Vero E6 Cells <sup>1</sup> Using RT-PCR <sup>3</sup>	Report results	1.1 × 10 <sup>4</sup> TCID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>5</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>Vero E6 cells; ATCC® CRL-1586™

<sup>2</sup>Grown in Eagle's Minimum Essential Medium (ATCC® 30-2003™) supplemented with 2% fetal bovine serum (ATCC® 30-2020™) for 14 days at 37°C and 5% CO<sub>2</sub>

<sup>3</sup>RNA used for RT-PCR was extracted from infected cell lysate and supernatant after 14 days incubation at 37°C and 5% CO<sub>2</sub>.

<sup>4</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) endpoint is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>5</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Date:** 18 MAR 2014

**Signature:** 

**Title:** Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

