SUPPORTING INFECTIOUS DISEASE RESEARCH

Salmonella enterica subsp. enterica, Strain 14028s ASTM14:2708 (Serovar Typhimurium)

Catalog No. NR-40615

For research use only. Not for human use.

Contributor:

Fred Heffron, Professor, Molecular Microbiology and Immunology Department, Oregon Health and Science University, Portland, Oregon, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: Enterobacteriaceae, Salmonella Species: Salmonella enterica Subspecies: Salmonella enterica subsp. enterica^{1,2} Serogroup: B Serovar: Typhimurium Strain: 14028s ΔSTM14:2708 (STM:2195)

- Original Source: Salmonella enterica (S. enterica) subsp. enterica, strain 14028s ΔSTM14:2708 (serovar Typhimurium) was derived from strain 14028s (strain 14028s was originally known as strain 14028, however, a variant of the original strain with a rough colony morphology was designated 14028r and the original smooth strain was renamed 14028s). Strain 14028 is a descendent of strain CDC 6516-60 which was isolated from pools of hearts and livers of 4-week-old chickens.⁴
- Comments: The ΔSTM14:2708 mutant of strain 14028s was produced by creating a PCR product with STM14:2708 homologous sequences at the 5' and 3' ends of a linear fragment containing a kanamycin resistance cassette. S. enterica subsp. enterica. strain 14028s was transformed. and insertion of the kan cassette in place of STM14:0807 was confirmed by PCR. The final non-polar deletions were constructed by elimination of the kan cassette.⁵ The complete genome (GenBank: CP001363.1) and plasmid (GenBank: CP001362.1) sequences are available for strain 14028s.

S. enterica are Gram-negative, rod-shaped, flagellated bacteria. The species is divided into six subspecies (I, II, IIIa, IIIb, IV, VI) where only subspecies I, subsp. enterica, is considered of clinical relevance. Salmonellosis (nontyphoidal), due to the greater than 1500 serovars of S. enterica subsp. enterica, is one of the most common foodborne diseases with an estimated two million cases that occur in the United States every year.⁶ Pathogenicity results from a variety of virulence factors found in plasmids, prophages, and five pathogenicity islands which allow these organisms to colonize and infect host organisms.

S. enterica subsp. enterica serovar Typhimurium (formerly

Salmonella typhimurium) is a major cause of gastroenteritis. These bacteria are host generalists that occur in humans and many other mammals. Septic shock resulting in part from lipopolysaccharide (LPS) is a primary complication associated with serovar Typhimurium infection.⁸ Due to its similarity to the clinical and pathological effects in humans, calves are currently used as an animal model for human enterocolitis caused by this serotype.9 Additionally, this serovar causes typhoid-like disease in mice and is used as a mouse model of human typhoid fever.4

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in 0.5X LB broth (supplemented with 1% Bacto-tryptone, 0.5% yeast extract and 1% sodium chloride) with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-40615 was packaged aseptically, in screw-capped plastic cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy broth or Nutrient broth or equivalent

Tryptic Soy agar with 5% defibrinated sheep blood or Nutrient agar or equivalent

Incubation:

Temperature: 37°C

Atmosphere: Aerobic

Propagation:

- Keep vial frozen until ready for use, then thaw. 1.
- Transfer the entire thawed aliquot into a single tube of 2. broth
- Use several drops of the suspension to inoculate an 3. agar slant and/or plate.
- 4. Incubate the tube, slant and/or plate at 37°C for 24 hours.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: Salmonella enterica subsp. enterica, Strain 14028s ΔSTM14:2708 (Serovar Typhimurium), NR-40615."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see www.cdc.gov/biosafety/publications/bmbl5/index.htm.

E-mail: contact@beiresources.org Tel: 800-359-7370 Fax: 703-365-2898

b|**e**|**i** resources

SUPPORTING INFECTIOUS DISEASE RESEARCH

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at <u>www.beiresources.org</u>.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC[®] nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC[®] nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC[®] and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC[®], their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, noncommercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

References:

- Judicial Commission of the International Committee on Systematics of Prokaryotes. "The Type Species of the Genus Salmonella Lignierres 1900 Is Salmonella enterica (ex Kauffmann and Edwards 1952) Le Minor and Popoff 1987, with the Type Strain LT2^T, and Conservation of the Epithet enterica in Salmonella enterica over All Earlier Epithets that May Be Applied to This Species. Opinion 80." <u>Int. J. Syst. Evol. Microbiol.</u> 55 (2005): 519-520. PubMed: 15653929.
- Tindall, B. J., et al. "Nomenclature and Taxonomy of the Genus Salmonella." <u>Int. J. Syst. Evol. Microbiol.</u> 55 (2005): 521-524. PubMed: 15653930.
- 3. F. Heffron, Personal Communication.
- Jarvik, T., et al. "Short-Term Signatures of Evolutionary Change in the Salmonella enterica Serovar Typhimurium 14028 Genome." <u>J. Bacteriol.</u> 192 (2010): 560-567. PubMed: 19897643.
- Datsenko, K. A. and B. L. Wanner. "One-Step Inactivation of Chromosomal Genes in *Escherichia coli* K-13 Using PCR Products." <u>Proc. Natl. Acad. Sci. USA</u> 97 (2000): 6640-6645. PubMed: 10829079.

- Altekruse, S. F., M. L. Cohen and D. L. Swerdlow. "Emerging Foodborne Diseases." <u>Emerg. Infect. Dis.</u> 3 (1997): 285-293. PubMed: 9284372.
- Lavigne, J. P. and A. B. Blanc-Potard. "Molecular Evolution of Salmonella enterica Serovar Typhimurium and Pathogenic Escherichia coli: From Pathogenesis to Therapeutics." <u>Infect. Genet. Evol.</u> 8 (2008): 217-226. PubMed: 18226587.
- Sha, J., et al. "The Two Murein Lipoproteins of Salmonella enterica Serovar Typhimurium Contribute to the Virulence of the Organism." <u>Infect. Immun.</u> 72 (2004): 3987-4003. PubMed: 15213144.
- Zhang, S., et al. "The Salmonella enterica Serotype Typhimurium Effector Proteins SipA, SopA, SopB, SopD, and SopE2 Act in Concert to Induce Diarrhea in Calves." <u>Infect. Immun.</u> 70 (2002): 3843-3855. PubMed: 12065528.

ATCC[®] is a trademark of the American Type Culture Collection.

