

Certificate of Analysis for NR-46412

Staphylococcus aureus, Strain HIP13170

Catalog No. NR-46412

Product Description: Staphylococcus aureus (S. aureus), strain HIP13170 was isolated in 2004 in New York, USA, from urine of a 63-year-old female with a polymicrobial infected nephrostomy tube exit site who had no recent history of vancomycin therapy. S. aureus, strain HIP13170 is a vancomycin-resistant S. aureus (VRSA) strain and is reported to be resistant to a number of other antimicrobial agents. It was co-isolated with S. aureus, strain HIP13419 which are similar in most aspects except the vancomycin resistant phenotype for HIP13170 is less stable than that of HIP13419

Lot¹: 62471638 Manufacturing Date: 03APR2014

TEST	SPECIFICATIONS	RESULTS
	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology	Gram-positive cocci	Gram-positive cocci
Colony morphology ²	Report results	Circular, convex, entire, smooth and
		cream (Figure 1)
Motility (wet mount)	Report results	Non-motile
Hemolysis ³	Report results	β-hemolytic
Biochemical Characterization		
Catalase	Positive	Positive
Coagulase ⁴	Report results	Positive
VITEK [®] 2 Compact (GP card)	Consistent with S. aureus	Consistent with S. aureus
VITEK® MS (MALDI-TOF)	Consistent with S. aureus	Consistent with S. aureus
Antibiotic Susceptibility Profile		
VITEK [®] (AST-GP ⁷ 1 card) ⁵		1
Beta-lactamase ⁶	Report results	Positive
Cefoxitin screen	Report results	Positive
Benzylpenicillin	Report results	Resistant (≥ 0.5 µg/mL)
Oxacillin	Resistant	Resistant (≥ 4 µg/mL)
Ciprofloxacin	Resistant	Resistant (≥ 8 µg/mL)
Levofloxacin	Report results	Resistant (≥ 8 µg/mL)
Moxifloxacin	Report results	Resistant (= 4 µg/mL)
Clindamycin (inducible resistance)	Report results	Negative
Erythromycin	Resistant	Resistant (≥ 8 µg/mL)
Clindamycin	Resistant	Resistant (≥ 8 µg/mL)
Quinupristin/dalfopristin	Sensitive	Sensitive (= 0.5 µg/mL)
Linezolid	Sensitive	Sensitive (= 0.5 µg/mL)
Daptomycin	Report results	Sensitive (= 2 µg/mL) Sensitive (= 0.25 µg/mL)
	Resistant	
Vancomycin		Resistant (≥ 32 μg/mL)
Minocycline	Report results	Sensitive (= 2 µg/mL)
Tetracycline	Report results	Resistant (≥ 16 µg/mL)
Tigecycline	Report results	Sensitive (≤ 0.12 μg/mL)
Nitrofurantoin	Report results	Sensitive (= 32 µg/mL)
Rifampicin	Report results	Sensitive (≤ 0.5 μg/mL)
Trimethoprim/sulfamethoxazole	Sensitive	Sensitive (≤ 10 μg/mL)
Etest [®] antibiotic test strips ⁷		
Chloramphenicol ⁸	Report results	Sensitive (= 3 µg/mL)
Teicoplanin ⁸	Report results	Resistant (= 48 μg/mL)
Gentamicin ⁸	Resistant	Resistant (= 16 μg/mL)

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898




SUPPORTING INFECTIOUS DISEASE RESEARCH

Certificate of Analysis for NR-46412

Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1490 base pairs)	Consistent with S. aureus	Consistent with S. aureus
Viability (post-freeze) ²	Growth	Growth

S. aureus, strain HIP13170 was deposited to BEI Resources as part of the NARSA collection. NR-46412 was produced by inoculation of the deposited material into Brain Heart Infusion broth with 6 µg/mL vancomycin and grown 25 hours at 37°C in an aerobic atmosphere. Broth inoculum was added to Brain Heart Infusion agar with 6 µg/mL vancomycin kolles which were grown 23 hours at 37°C in an aerobic atmosphere to produce this lot. Purity of this lot was assessed for 7 days under propagation conditions.

Figure 1



Date: 27 JUN 2014

Signature: (

Title:

Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²21 hours at 37°C in an aerobic atmosphere on Brain Heart Infusion agar with 6 μg/mL vancomycin

³21 hours at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood

⁴4 hours at 37°C in rabbit serum with 0.15% EDTA (Coagulase Plasma BBL™ 240827)

⁵Minimum Inhibitory Concentration (MIC); MIC Interpretation Guideline: CLSI M100-S22 (2012)

⁶The production of beta-lactamase was detected using a Cefinase™ Paper Disc (BBL™ 231650)

⁷24 hours at 37°C in an aerobic atmosphere on Mueller Hinton agar

⁸For both chloramphenicol (bioMérieux Etest[®] 412308) and teicoplanin (bioMérieux Etest[®] 412459), a MIC ≤ 8 μg/mL is sensitive, a MIC = 16 μg/mL is intermediate and a MIC ≥ 32 μg/mL is resistant. For gentamicin (bioMérieux Etest[®] 412367), a MIC ≤ 4 μg/mL is sensitive, a MIC = 8 μg/mL is intermediate and a MIC ≥ 16 μg/mL is resistant.