

Human Metapneumovirus TN/91-316

Catalog No. NR-22232

Product Description: Cell lysate and supernatant from *Macaca mulatta* kidney epithelial cells¹ infected with human metapneumovirus (HMPV), TN/91-316

Passage History: L8/L5 (Vanderbilt/BEI Resources; L# = Number of passages in LLC-MK2 cells)

Lot²: 61746424

Manufacturing Date: 16DEC2013

| TEST | SPECIFICATIONS | RESULTS |
|--|---|---|
| Identification by Infectivity in LLC-MK2 Derivative Cells ¹ | Report results | Rounding and dissociation from the monolayer |
| Genome Sequencing ³ | Consistent with HMPV TN/91-316 | 99% identity with HMPV TN/91-316 (GenBank: KC403971) |
| Titer by TCID ₅₀ Assay ^{4,5} in LLC-MK2 Derivative Cells ¹ With Direct Fluorescence Assay (DFA) Readout ⁶ | Report results | 1.0 x 10 ⁶ TCID ₅₀ per mL |
| Sterility (21-day incubation) Harpo's HTYE broth ⁷ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂ | No growth No growth No growth No growth No growth No growth No growth | No growth No growth No growth No growth No growth No growth No growth |
| Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid | None detected None detected | None detected None detected |

¹LLC-MK2 Derivative cells (ATCC® CCL-7.1™)

²Grown in Opti-MEM® Minimal Essential Medium (Life Technologies 31985) supplemented with 2 mM L-glutamine (Life Technologies 25030), 100 µg per mL CaCl₂, and 5 µg per mL trypsin (Gibco® 25200-056) for 7 days at 37°C and 5% CO₂

³The nucleotide sequencing of NR-22232 performed at BEI Resources cannot confirm exact strain identity owing to the high degree of sequence conservation within HMPV lineages.

⁴The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁵13 days at 37°C and 5% CO₂

⁶Using Light Diagnostics™ Human Metapneumovirus DFA Reagent (Millipore 3124)

⁷Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 29 OCT 2014

Signature: 

Title: Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

