

## Certificate of Analysis for NR-44354

## Klebsiella pneumoniae, Strain UHKPC45

## Catalog No. NR-44354


**Product Description:** *Klebsiella pneumoniae* (*K. pneumoniae*), strain UHKPC45 was isolated in 2008 from blood of a patient with sepsis in Cleveland, Ohio, USA.

Lot<sup>1</sup>: 62849224 Manufacturing Date: 14AUG2014

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology Colony morphologies <sup>2,3</sup>	Gram-negative bacilli	Gram-negative bacilli
Colony morphologies <sup>2,3</sup>	Report results	Colony type 1: Circular, low convex, entire, smooth, mucoid and gray (Figure 1)
		Colony type 2: Circular, convex, entire, smooth, mucoid and light gray (Figure 1)
Motility (wet mount)	Report results	Non-motile
VITEK®MS (MALDI-TOF)	Consistent with K. pneumoniae	Consistent with K. pneumoniae
Genotypic Analysis		
Sequencing of 16S ribosomal RNA gene (~ 1470 base pairs) <sup>3</sup>	Consistent with K. pneumoniae	Consistent with K. pneumoniae
Riboprinter® Microbial Characterization System	Consistent with K. pneumoniae	Consistent with K. pneumoniae
Purity (post-freeze) <sup>4</sup>	Consistent with K. pneumoniae	Consistent with K. pneumoniae
Viability (post-freeze) <sup>2</sup>	Growth	Growth

<sup>&</sup>lt;sup>1</sup>K. pneumoniae, strain UHKPC45 was deposited by Robert A. Bonomo, M.D., Chief of Medical Services, Louis Stokes Cleveland VA Medical Center, Cleveland, Ohio, USA. NR-44354 was produced by inoculation of the deposited material into Tryptic Soy broth and incubated for 24 hours at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar with 5% defibrinated sheep blood kolles which were grown 22 hours at 37°C in an aerobic atmosphere to produce this lot.

<sup>&</sup>lt;sup>4</sup>Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood.



BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

<sup>&</sup>lt;sup>2</sup>22 hours at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood.

<sup>&</sup>lt;sup>3</sup>Two colony types were observed. Plating of the individual colony types showed that they did not revert to a single colony type. The 16S ribosomal RNA gene of each colony type was sequenced and found to be consistent with the other colony type and with *K. pneumoniae*.



## Certificate of Analysis for NR-44354

**Date:** 11 DEC 2014

Signature:

**BEI Resources Authentication** 

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

 $\mathsf{ATCC}^{\$}$  is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898