

***Fusobacterium nucleatum*, Strain CTI-02**

**Catalog No. HM-993**

**Product Description:** *Fusobacterium nucleatum* (*F. nucleatum*), strain CTI-02 was isolated in 2012 from colonic tumor tissue from a human patient with colorectal carcinoma in Massachusetts, USA.

**Lot<sup>1,2</sup>: 63140958**

**Manufacturing Date: 16JAN2015**

TEST	SPECIFICATIONS	RESULTS
<b>Phenotypic Analysis</b> Cellular morphology Colony morphology <sup>3</sup>  Motility (wet mount)	Gram-negative rods Report results  Report results	Gram-negative rods Circular, convex, entire, opaque, smooth and gray (Figure 1) Non-motile
<b>Genotypic Analysis</b> Sequencing of 16S ribosomal RNA gene (~ 1410 base pairs)	≥ 99% identical to GenBank: AXNY01000032 ( <i>F. nucleatum</i> , strain CTI-02)	≥ 99% identical to GenBank: AXNY01000032 ( <i>F. nucleatum</i> , strain CTI-02)
<b>Purity (post-freeze)</b> Anaerobic growth <sup>4</sup> Aerobic growth <sup>5</sup>	Growth consistent with <i>F. nucleatum</i> No growth	Growth consistent with <i>F. nucleatum</i> No growth
<b>Viability (post-freeze)<sup>3</sup></b>	Growth	Growth

<sup>1</sup>Quality control of HMP material is only performed to demonstrate that the material distributed by BEI Resources is identical to the deposited material. It should not be considered a complete characterization of the deposited organism.


<sup>2</sup>*F. nucleatum*, strain CTI-02 was deposited by Wendy S. Garrett, M.D., Ph.D., Assistant Professor, and Aleksander D. Kostic, Department of Immunology and Infectious Diseases, Harvard School of Public Health, Boston, Massachusetts, USA. HM-993 was produced by inoculation of the deposited material into Modified Chopped Meat medium and incubated for 47 hours at 37°C in an anaerobic atmosphere (< 5% O<sub>2</sub>; Remel™ Pack-Anaero™). Broth inoculum was added to Tryptic Soy agar with 5% defibrinated sheep blood kolles which were grown 46 hours at 37°C in an anaerobic atmosphere to produce this lot.

<sup>3</sup>48 hours at 37°C in an anaerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood

<sup>4</sup>Purity of this lot was assessed for 7 days on Tryptic Soy agar with 5% defibrinated sheep blood at 37°C in an anaerobic atmosphere.

<sup>5</sup>Purity of this lot was assessed for 48 hours on Tryptic Soy agar with 5% defibrinated sheep blood at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub>.

**Figure 1**



**Date:** 16 MAR 2015

**Signature:**

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

