

Influenza A Virus, A/California/04/2009 (H1N1)pdm09, Cell Isolate (Produced in Eggs)

Catalog No. NR-13659

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs¹ infected with influenza A virus, A/California/04/2009 (H1N1)pdm09.

Passage History: C2/E2 (CDC/BEI); C# = Number passages in MDCK cells, E# = Number of passages in embryonated chicken eggs

Lot²: 63633075

Manufacturing Date: 29JUL2015

TEST	SPECIFICATIONS	RESULTS
Sequencing of Hemagglutinin and Matrix Coding Regions Hemagglutinin (495 nucleotides) Matrix (944 nucleotides)	Consistent with A/California/04/2009 (H1N1)pdm09 Consistent with A/California/04/2009 (H1N1)pdm09	99% identity with A/California/04/2009 (H1N1)pdm09 (GenBank: JF915184) 99% identity with A/California/04/2009 (H1N1)pdm09 (GenBank: FJ969513)
Titer by CEID₅₀ Assay^{3,4} in Embryonated Chicken Eggs¹	Report results	2.8 × 10 ⁷ CEID ₅₀ per mL
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Brucella agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹9 day old SPF Embryonated Chicken Eggs acquired from B&E Eggs, York Springs, Pennsylvania

²Grown in the allantoic cavity of embryonated chicken eggs¹ for 2 days at 33.5°C in a humidified chamber without CO₂

³The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁴2 days at 33.5°C in a humidified chamber

⁵Atlas, Ronald M. Handbook of Microbiological Media. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 11 NOV 2015

Signature: 

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

