

**Zika Virus, MR 766**

**Catalog No. NR-50065**

**Product Description:** Cell lysate and supernatant from *Cercopithecus aethiops* kidney epithelial cells (Vero)<sup>1</sup> infected with Zika virus (ZIKV), MR 766

**Passage History:** SM150/V8 (Prior to deposit at BEI/BEI); SM# = Number of passages in suckling mice; V# = Number of passages in Vero cells

**Lot<sup>2</sup>: 63856750**

**Manufacturing Date: 18JAN2016**

TEST	SPECIFICATIONS	RESULTS
Infectivity in Vero Cells <sup>1</sup>	Report results	Cell rounding and detachment
Sequencing of Species-Specific Region (887 nucleotides)	Consistent with ZIKV, MR 766	100% identity with ZIKV, MR 766 (GenBank: AY632535) <sup>3</sup>
Titer by TCID <sub>50</sub> Assay in Vero Cells <sup>1,4,5</sup>	Report results	1.6 × 10 <sup>8</sup> TCID <sub>50</sub> per mL
<b>Sterility (21-day incubation)</b> Harpo's HTYE broth <sup>6</sup> , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO <sub>2</sub>	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
<b>Mycoplasma Contamination</b> Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

<sup>1</sup>Vero cells: ATCC® CCL-81™

<sup>2</sup>Grown in Dulbecco's Modified Eagle's Medium containing 4 mM L-glutamine, 4500 mg per L glucose, 1 mM sodium pyruvate, and 1500 mg per L sodium bicarbonate (ATCC® 30-2002), supplemented with 2% fetal bovine serum (ATCC® 30-2020) for 2 days at 37°C with 5% CO<sub>2</sub>

<sup>3</sup>The complete coding sequence of NR-50065, Lot No. 63856750 has also been determined (GenBank: KU963573).

<sup>4</sup>The Tissue Culture Infectious Dose 50% (TCID<sub>50</sub>) is the 50% infectious endpoint in cell culture. The TCID<sub>50</sub> is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD<sub>50</sub>) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID<sub>50</sub> provides a measure of the titer (or infectivity) of a virus preparation.

<sup>5</sup>8 days at 37°C with 5% CO<sub>2</sub>

<sup>6</sup>Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

**Date:** 06 APR 2016

**Signature:**



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

