

Certificate of Analysis for NR-15772

Kaeng Khoi Virus, PSC-19

Catalog No. NR-15772

Product Description: Clarified supernatant from *Cercopithecus aethiops* kidney epithelial cells (Vero E6)¹ infected with Kaeng Khoi virus (KKV), PSC-19

Lot²: 59029796 Manufacturing Date: 2009

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity in Vero E6 Cells ¹	Report results	Cell rounding and sloughing
Sequencing of Segment L Polymerase Gene (954 nucleotides)	Consistent with KKV, PSC-19	99% identity with KKV, PSC-19 (GenBank: KJ867205)
Titer by TCID ₅₀ Assay ³ in Vero E6 Cells ^{1,4}	Report results	$2.8 \times 10^6 \text{ TCID}_{50} \text{ per mL}$
Functional Activity by RT-PCR Assay	~1200 bp amplicon	~1200 bp amplicon
Sterility (21-day incubation) Harpo's HTYE broth ⁵ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Sheep blood agar, 37°C, aerobic Sheep blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth	No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA Detection by PCR of Test Article nucleic acid	None detected None detected	None detected None detected

¹Vero E6 cells: ATCC[®] CRL-1586™

Date: 25 APR 2016

ignature:

BEI Resources Authentication

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

BEI Resources www.beiresources.org E-mail: contact@beiresources.org
Tel: 800-359-7370

Fax: 703-365-2898

²Grown and deposited by Charles H. Calisher, Ph.D., Department of Microbiology, Immunology and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, Colorado, USA

³The Tissue Culture Infectious Dose 50% (TCID₅₀) endpoint is the 50% infectious endpoint in cell culture. The TCID₅₀ is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the culture vessels inoculated, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the TCID₅₀ provides a measure of the titer (or infectivity) of a virus preparation.

⁴8 days at 37°C and 5% CO₂

⁵Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.