

Staphylococcus aureus, Strain NY-177

Catalog No. NR-46237

Product Description: *Staphylococcus aureus* (*S. aureus*), strain NY-177 was isolated in 2005 from a bursa of a 90-year-old male with bursitis and/or cellulitis in New York, USA. *S. aureus*, strain NY-177 is a methicillin-resistant *S. aureus* (MRSA) strain.

Lot¹: 64044911

Manufacturing Date: 18FEB2016

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ² Motility (wet mount) Hemolysis ² Biochemical characterization Catalase Coagulase ³ VITEK [®] 2 Compact (GP card) VITEK [®] MS (MALDI-TOF)	Gram-positive cocci Report results Report results Report results Positive Report results ≥ 90% probability of being <i>S. aureus</i> <i>S. aureus</i>	Gram-positive cocci Circular, convex, entire, smooth and cream (Figure 1) Non-motile β-hemolytic Positive Positive <i>S. aureus</i> (99% probability) ⁴ <i>S. aureus</i> (99.9%)
Antibiotic Susceptibility Profile VITEK [®] (AST-GP71 card) ⁵ Beta-lactamase ⁶ Cefoxitin screen Benzylpenicillin Oxacillin Ciprofloxacin Levofloxacin Moxifloxacin Clindamycin (inducible resistance) Erythromycin Clindamycin Quinupristin/dalfopristin Linezolid Daptomycin Vancomycin Minocycline Tetracycline Tigecycline Nitrofurantoin Rifampicin Trimethoprim/sulfamethoxazole Etest [®] antibiotic test strips ⁸ Chloramphenicol ⁹ Teicoplanin ⁹ Gentamicin ⁹	Report results Report results Report results Resistant Report results Resistant Report results Report results Resistant Resistant Report results Sensitive Sensitive Sensitive Report results Resistant Report results Report results Sensitive Resistant Report results Report results Resistant	Positive Positive Resistant (≥ 0.5 µg/mL) Resistant (≥ 4 µg/mL) Resistant (≥ 8 µg/mL) Resistant (≥ 8 µg/mL) Resistant (4 µg/mL) Negative Resistant (≥ 8 µg/mL) Resistant (≥ 8 µg/mL) Sensitive (0.5 µg/mL) Sensitive (2 µg/mL) Sensitive (0.5 µg/mL) Sensitive (≤ 0.5 µg/mL) Sensitive (4 µg/mL) Resistant (≥ 16 µg/mL) Sensitive (≤ 0.12 µg/mL) ⁷ Sensitive (≤ 16 µg/mL) Sensitive (≤ 0.5 µg/mL) Resistant (≥ 320 µg/mL) Sensitive (8 µg/mL) ¹⁰ Sensitive (1.5 µg/mL) Resistant (≥ 256 µg/mL) ¹¹
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (~ 1480 base pairs)	≥ 99% sequence identity to <i>S. aureus</i> type strain (GenBank: L37597)	100% sequence identity to <i>S. aureus</i> type strain (GenBank: L37597)
Purity (post-freeze)¹²	Consistent with expected colony morphology	Consistent with expected colony morphology
Viability (post-freeze)²	Growth	Growth

- ¹*S. aureus*, strain NY-177 was deposited to BEI Resources as part of the NARSA collection. NR-46237 was produced by inoculation of the deposited material into Tryptic Soy broth and grown 1 day at 37°C in an aerobic atmosphere. Broth inoculum was added to Tryptic Soy agar with 5% defibrinated sheep blood kolles which were grown 1 day at 37°C in an aerobic atmosphere to produce this lot.
- ²1 day at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood
- ³4 hours at 37°C in rabbit serum with 0.15% EDTA (Coagulase Plasma BBL™ 240827)
- ⁴Percent probabilities above 90% indicate a close match to the typical biochemical pattern for the given organism, with a percent probability of 99% being a perfect match between the test reaction pattern and the unique biochemical pattern of the given organism or organism group. For additional information, please refer to O'Hara, C.M. and J. M. Miller. "Evaluation of the Vitek 2 ID-GNB Assay for Identification of Members of the Family *Enterobacteriaceae* and Other Nonenteric Gram-Negative Bacilli and Comparison with the Vitek GNI+ Card." *J. Clin. Microbiol.* 41 (2003): 2096-2101. PubMed: 12734254.
- ⁵Minimum Inhibitory Concentration (MIC); MIC Interpretation Guideline: CLSI M100-S22 (2012)
- ⁶The production of beta-lactamase was detected using a Cefinase™ Paper Disc (BBL™ 231650).
- ⁷MIC Interpretation Guideline: EUCAST Version 4.0 (2014)
- ⁸1 day at 37°C in an aerobic atmosphere on Mueller Hinton agar
- ⁹For both chloramphenicol (bioMérieux Etest® 412308) and teicoplanin (bioMérieux Etest® 412459), a MIC ≤ 8 µg/mL is sensitive, a MIC = 16 µg/mL is intermediate and a MIC ≥ 32 µg/mL is resistant. For gentamicin (bioMérieux Etest® 412367), a MIC ≤ 4 µg/mL is sensitive, a MIC = 8 µg/mL is intermediate and a MIC ≥ 16 µg/mL is resistant.
- ¹⁰*S. aureus*, strain NY-177 was deposited as having an intermediate susceptibility to chloramphenicol. Antibiotic susceptibility testing performed in duplicate identified strain NY-177 as being susceptible to chloramphenicol.
- ¹¹Susceptibility testing using gentamicin Etest® stripes resulted in an inhibition ellipse consistent with a susceptible to intermediately susceptible phenotype. However, within the ellipse, there were satellite cells that were highly resistant to gentamicin, suggesting that *S. aureus*, strain NY-177 is composed of a mixed population of cells and should be considered resistant to gentamicin.
- ¹²Purity of this lot was assessed for 7 days at 37°C in an aerobic atmosphere on Tryptic Soy agar with 5% defibrinated sheep blood.

Figure 1: Colony Morphology



Date: 11 JUL 2016

Signature:

BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

